



HAL
open science

Moringa Oleifera supplemented biscuits: Nutritional values and consumer segmentation

Amel Hedhili, Samuel Lubbers, Elias Bou-Maroun, Francine Griffon, Babatope Ebenezer Akinyemi, Florence Husson, Dominique Valentin

► To cite this version:

Amel Hedhili, Samuel Lubbers, Elias Bou-Maroun, Francine Griffon, Babatope Ebenezer Akinyemi, et al.. Moringa Oleifera supplemented biscuits: Nutritional values and consumer segmentation. South African Journal of Botany, 2021, 138, pp.406-414. 10.1016/j.sajb.2021.01.017 . hal-03187745

HAL Id: hal-03187745

<https://institut-agro-dijon.hal.science/hal-03187745v1>

Submitted on 13 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 ***Moringa Oleifera* supplemented biscuits: nutritional values and consumer**
2 **segmentation**

3 Amel Hedhili^{a,b,*}, Samuel Lubbers^b, Elias Bou-Maroun^b, Francine Griffon^a, Babatope
4 Ebenezer Akinyemi^c, Florence Husson^b, Dominique Valentin^a

5 ^a: Centre des Sciences du Goût et de l'Alimentation, AgroSup Dijon, CNRS, INRAE,
6 Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

7 ^b: Univ. Bourgogne Franche-Comté, AgroSup Dijon, PAM UMR A 02.102, F-21000
8 Dijon, France

9 ^c: Department of Agricultural Economics & Extension. University of Fort Hare, Alice,
10 5700, South Africa

11 *Corresponding author. E-mail address: Amel.Hedhili@inrae.fr (A. Hedhili)

12

13 **Abstract**

14 *Moringa oleifera* is a nutrient-rich plant, known as a miracle tree, and commonly used
15 as a food supplement. Despite its popularity, there are scant empirical findings on its
16 real nutritional benefits and acceptance among consumers when added to food. To
17 fill this gap, we added Moringa Dried Leaves (MDL) to biscuits. A consumer test was
18 implemented to study the sensory properties of the biscuits. Nutritional properties,
19 including in vitro protein digestibility, and physical properties were also investigated.
20 Global liking indicates the segmentation of consumers into two clusters and
21 confirmed heterogeneity. Cluster 1 consumers disliked the supplemented biscuits
22 and cited dark green color and texture as the main reasons for dislike. Conversely,
23 cluster 2 consumers, who were much younger, liked the supplemented biscuits due
24 to their adventurous taste for novel products. Nutritional analysis revealed that the
25 incorporation of MDL to biscuits increased their iron and proteins but led to
26 incomplete protein digestion especially for biscuits with high supplementation. In
27 addition, vitamin A derived from MDL was degraded during baking. The biscuit with
28 the highest acceptability and digestion (C1) would increase protein intake only slightly
29 but provide essential amino acids, which are deficient in wheat flour. These findings,
30 confirm that there is a limit to further increasing the amount of MDL added to foods in
31 order to provide more nutrients. In addition, it is recommended that Moringa
32 supplemented products should be advertised with caution in order not to mislead the
33 consumers.

34 **Keywords:** Moringa Dried Leaves (MDL); Hedonic test; Consumer segmentation;
35 Open-ended questions; In vitro digestibility; Proximate analysis

36 1. Introduction

37 The *Moringa oleifera* tree is universally known as a miracle plant. This plant derived
38 its name from the fact that all its parts (leaves, seeds, flowers etc.) can be used as
39 nutritional supplement, medicine, water purifier and animal fodder (Daba, 2016).
40 Native to northern India, *Moringa oleifera* Lam. (*Moringaceae*) is now cultivated in
41 many subtropical regions in Africa, tropical America, Mexico, Malaysia, and the
42 Philippines (Adeyemi et al., 2014; Fahey, 2005). Several studies have shown that
43 *Moringa oleifera* leaves are a rich source of certain macro and micronutrients.
44 Sánchez-Machado et al. (2010) reported that Moringa plant contains a high
45 percentage of essential amino acids in green leaves. While Siddhuraju and Becker
46 (2003) and Sreelatha and Padma (2009) have also reported antioxidant properties of
47 *Moringa oleifera* leaves. The leaves of Moringa have also been found to contain
48 important amounts of calcium, potassium, magnesium, vitamin A, C and E (Hekmat,
49 2015). Witt's (2008) meta-analysis of 49 scientific articles summarizes the nutritional
50 composition of Moringa dried leaves (MDL). It shows that in 100 g of MDL there are
51 24 ± 5.8 g of proteins, 36 ± 9.2 g of carbohydrates, 6 ± 2.5 g of fat, 20.6 to 28.6 g of
52 fiber, 32.5 ± 10.8 mg of iron, and 3639 ± 1980 μ g of vitamin A.

53 Currently, the use of *Moringa oleifera* as a nutrient-rich plant for food
54 supplementation is getting much attention. It was first used in India before it was later
55 exported to African countries where it was incorporated in traditional foods such as
56 amala, a starchy meal in Nigeria and Ghana (Oyeyinka and Oyeyinka, 2016). In
57 developed countries, for instance in Europe, *Moringa oleifera* is marketed as a
58 “superfood” for a wide range of health benefits such as high level of antioxidant
59 activity (Bruins et al., 2018). The importation of Moringa to Europe was estimated

60 between 100 and 200 tons in 2017 and the main market destinations are the United
61 Kingdom, Germany, and France (CBI, 2019).

62 Many scientific studies have shown that the supplementation of staple foods such as
63 bread, cheese, yoghurt, and biscuits with MDL improve their nutritional properties
64 (Dachana et al., 2010; Hekmat et al., 2015; Salem et al., 2013; Sengeev et al., 2013).

65 Biscuits is one of the most popular bakery products in the human diet due to many
66 characteristics such as palatability, convenience, ready-to-eat, affordability and
67 longer shelf-life in comparison to the other bakery products (Arshad, 2007; Nagi,
68 2012). It is also widely accepted and consumed in many countries, which makes it a
69 viable supplementation vehicle for nutritional improvement of vital nutrients like
70 protein, dietary fibers and functional ingredients (Sulieman et al., 2019).

71 In previous studies, MDL were incorporated in biscuits with levels varying from 3% to
72 15% and results showed that this supplementation increased protein, fiber, b-
73 carotene (Dachana et al., 2010; Mouminah, 2019), and minerals (Fe, Ca, Mg, P and
74 K) composition of biscuits (Abdel-Samie and Abdulla, 2014). The incorporation of
75 Moringa also has an impact on the physical and sensory properties of biscuits.

76 Physical properties such as diameter and thickness changed depending on the
77 amount of MDL added and breaking force significantly increased with the
78 concentration of Moringa (Dachana et al., 2010). Previous studies that focused on
79 sensory evaluation showed that biscuits were acceptable up to 9% or 10% of
80 Moringa supplementation respectively (Abdel-Samie and Abdulla, 2014; Dachana et
81 al., 2010). Lee et al. (2014) revealed that the typical leafy flavor and slightly bitter
82 taste decreased the consumer acceptance of a product supplemented with leaves.
83 Furthermore, Dachana et al. (2010) observed that the addition of MDL to cookies
84 resulted in darker samples, which can lead to a lower consumer acceptance. This

85 finding is in agreement with Ramota et al. (2013) who state that MDL has a deep
86 green color, which may be attributed to its high chlorophyll content and may strongly
87 mask the color of original foods.

88 In contrast, a study carried out by Ramaroson et al. (2014) in Madagascar showed
89 that the Moringa green color is not always a problem when added to food. Cassava
90 root cakes were supplemented with MDL and presented to children who had to score
91 their overall liking of the samples. At the end of the experience, children chose one of
92 the products as a reward for their participation. Results showed that the product
93 containing the highest proportion of Moringa received high liking score and was
94 chosen as reward by a majority of the children despite its greener color, which
95 demonstrated that color was not a barrier to children acceptance in the study. This
96 result suggests that there might be inter-individual difference between consumers in
97 terms of acceptance.

98 Previous studies investigated consumer preferences without examining the
99 underlining heterogeneity that might exist among consumers due to individual
100 peculiarity (Abdel-Samie and Abdulla, 2014; Dachana et al., 2010; Rathnayake and
101 Navarathna, 2017). Hence, the first objective of this study was to implement sensory
102 analysis in order to gain understanding of heterogeneity in consumer preferences of
103 Moringa supplemented biscuits by looking at inter-individual differences among the
104 study participants.

105 Another potential limitation of the use of Moringa for food supplementation is that
106 most of the reported evidence do not take into account the effect of processing on
107 nutritional compounds and/or of the bio-availability of these compounds.

108 Furthermore, the few studies which explored this last point are far from being
109 consensual. For example, a study carried out by Teixeira et al (2014), showed that

110 the in vitro protein digestibility (IVPD) of MDL was 33.29% whereas a previous study
111 comparing the IVPD in thirteen plants species reported a value of 74.2% of IVPD of
112 MDL (Ly et al., 2001). The second goal of this study was to investigate the level of
113 nutritional improvement of biscuits due to Moringa supplementation.

114 To achieve these objectives, our study asked and offered answers to four questions,
115 namely: (1) Did the incorporation of MDL in biscuits improve their nutritional
116 properties and affect their IVPD? (2) What is the contribution of biscuits containing
117 MDL in recommended daily requirement? (3) Did the incorporation of MDL in biscuits
118 affect the physical properties of biscuits? (4) What is the effect of supplementation
119 with MDL on sensory properties of biscuits and consumer acceptance?

120 **2. Materials and methods**

121 *2.1. Biscuit preparation*

122 All ingredients (Wheat flour T-45, sugar, shortening, salt and baking powder) used to
123 produce the biscuits were purchased from local supermarket Carrefour®, Dijon,
124 France. MDL were purchased in Dijon, France.

125 Three experimental biscuits (C1, C2, and C3) were prepared by supplementing a
126 control biscuit (C0) with different amounts of MDL. The control biscuit was made of
127 100 g wheat flour, 30 g sugar powder, 30 g shortening, 2.5 g baking powder, 0.5 g
128 salt, and water. A proportion of 10%, 20% and 30% of MDL, respectively was added
129 to the base recipe to obtain supplemented biscuits C1, C2 and C3, respectively.

130 Wheat flour and MDL were sieved before mixing with shortening until homogeneous
131 lumps were formed. Then, sugar, salt and baking powder were added. The mixture
132 was mixed at low speed with a pastry robot for 10 min with a stop every 2 min. Water

133 was added gradually and mixed for 2 min at low speed. The dough was wrapped in
134 stretch film and refrigerated at 4 °C for 20 min. After 20 minutes, the dough was
135 placed on a lightly floured biscuit sheet. The prepared dough was rolled to a
136 thickness of 4mm and cut using a circular shape of 3 cm in diameter. The biscuits
137 were baked at 180 °C for 20 min. After baking, the biscuits were cooled down to
138 room temperature and kept in airtight food plastic boxes. Fig. 1 shows the control
139 biscuit (C0) and the biscuits supplemented with the three levels of MDL (C1 to C3).

140 *2.2. Proximate analysis*

141 All reagents and chemicals used for the proximate analysis were purchased from
142 Sigma Aldrich® (France) and were of analytical grade. Biscuit samples were
143 analyzed in triplicate for moisture, proteins, vitamin A and iron. Moisture content was
144 determined according to AOAC (2011). Crude proteins were determined according to
145 *Kjeldahl* method. The proteins content was obtained using the conversion factor 4.4
146 (factor used for tropical plant samples) (Ramaroson et al., 2014). Vitamin A content
147 was determined using the method of vitamin A (Retinol) in foods (AOAC, 2001). Iron
148 content was obtained following the method of determination of trace elements by
149 atomic absorption spectrometry (French standard 14082, 2003).

150 *2.3. In vitro protein digestibility (IVPD)*

151 Pepsin (from porcine stomach mucosa, >2500 U/mg protein), pancreatin (from
152 porcine pancreas, 4 x USP), casein sodium salt from bovine milk (C-8654) and TAME
153 (p-Toluene-Sulfonyl-L-arginine methyl ester; T4626) were purchased from Sigma-
154 Aldrich (France). In vitro digestibility was carried out according to the method of
155 Minekus et al. (2014) with some modifications. Analysis was done in duplicate. In
156 addition, 1 g of MDL or 1 g of biscuit was prepared in 10 mL of solution then 8 mL of
157 simulated gastric fluid (SGF) electrolyte solution was added. To this mixture 1.6 mL

158 of porcine pepsin solution (25.000 U/mL) and 5 μ L of 0.3 M CaCl₂ were added. The
159 pH of the mixture was adjusted to 3.0. The solution was incubated for 120 min at 37
160 °C under continuous stirring at 170 rpm. After gastric digestion, gastric chime was
161 mixed with 11 mL of simulated intestinal fluid (SIF). To this solution, 736 mg of the
162 pancreatic enzyme and 40 μ L of 0.3 M CaCl₂ were added and the final volume was
163 completed to 40 mL by SIF solution. The pH of the mixture was adjusted to 7.0. The
164 final solution was incubated for 120 min at 37 °C with stirring. To determine the
165 nitrogen release in the samples, the trichloroacetic acid (TCA) precipitation method
166 was used. Aliquots of 10 mL of the withdrawn digestion mixture were mixed with 10
167 mL of 10% TCA solution. The mixtures were mixed and then centrifuged (8000 g, 10
168 min). The nitrogen content of the resulting supernatants was determined by the
169 *Kjeldahl* method. The percentage of soluble nitrogen released during the digestion
170 process was calculated using the following equation (Teixeira et al., 2014).

$$171 \quad \% \text{ Enzymatic Hydrolysis (\%EH)} = \frac{\text{amount of protein after digestion}}{\text{total amount of protein}} * 100$$

172 The percentage of digestibility of the proteins was deduced from the digestibility of
173 the casein considered as standard (100%) (Teixeira et al., 2014).

174 *2.4. Physical characteristics of biscuits*

175 *2.4.1. Diameter, thickness and spread ratio*

176 Diameter (mm), thickness (mm) and spread ratio of samples were measured in
177 triplicate following the method of Ganorkar and Jain (2014). To determine biscuit
178 diameter, six biscuits were placed edge to edge and the total diameter was measured
179 in mm by a Vernier caliper. Then biscuits were rotated at an angle 90° for triplicate
180 measure. The average diameter was reported in mm. Six biscuits were placed on top

181 of one another to determine thickness. The total height was measured and the
182 average was presented in mm. Spread ratio was calculated as: spread ratio =
183 Average diameter / Average thickness * 10.

184 *2.4.2. Breaking force*

185 A three-point bend test was used to estimate the biscuits breaking force as an
186 indicator of hardness using a texture analyzer TA-TX2 equipped with a 25 kg load
187 cell. The peak breaking force (N) of biscuits using the force in compression was
188 recorded. Biscuit samples were placed on base beams with a distance of 1.8 cm
189 between the two beams. A three point-bending rig was used with an HDP/3PB, knife-
190 edge probe. The analyzer was set at a return-to-start cycle with a pretest speed of 2
191 mm/s, a test speed of 2 mm/s, and a post-test speed of 10 mm/s. The trigger was 20
192 g and distance of 10 mm. Analysis was done in triplicate.

193 *2.4.3 Color measurement*

194 Color attributes of biscuits supplemented with MDL (L*, a* and b*) were measured in
195 triplicate using Hunter Lab color analyzer (HunterLab Color Flex EZ, USA). The L*
196 value (lightness index scale) ranges from 0 (black) to 100 (white) while, a* value
197 indicates the redness (+a*) or greenness (-a*) and the b* value refers to the
198 yellowness (+b*) or blue (-b*). The petri dish containing sample was placed directly
199 on the colorimeter sensor.

200 *2.5. Sensory evaluation*

201 *2.5.1. Consumers*

202 The consumer panel consisted of 91 students and staff members from the University
203 of Burgundy (73.63% Women – 26.37% Men; 74.63% <30 years and 25.27% >30

204 years). To ensure a larger generalizability of the results no constraint was placed on
205 the nationality of consumers: 75.82% were French, 10.99% were African, 5.49%
206 were European other than French and 7.69% Asian.

207 *2.5.2. Experimental conditions*

208 Consumers were provided with information about the study, following which they
209 completed an informed consent form in keeping with French ethical requirements.

210 The consumers tasted the samples in a sensory evaluation room at Agrosup Dijon.

211 The biscuits were served in covered white plastic cups coded with a three-digit
212 random number. Two series of four samples (C0, C1, C2, and C3) each were
213 presented on a tray. Different random codes and different presentation orders were
214 used for the two series. The presentation order followed Williams' Latin squares.

215 Consumers were not aware that the biscuits in the two series were the same. Bottled
216 water was provided for palate cleansing. The questionnaire was written in French
217 language and explained in English when necessary.

218 *2.5.3. Procedure*

219 The consumer test included two steps. During the first step, consumers were
220 instructed to taste the first series of samples from left to right. They had to rate the
221 overall liking of each sample on a structured scale ranging from 1 (I do not like at all)
222 to 7 (I like it a lot) with 4 being anchored as neither like nor dislike. Before proceeding
223 to the next samples, consumers were asked to write down what they liked and what
224 they disliked about the sample (Symoneaux and Galmarini, 2014). During the second
225 step, consumers were then asked to taste the second series of biscuits and evaluate
226 the appearance, odor, taste and texture of the samples using 7-points hedonic
227 scales. Consumers could answer either in French or in English as they wished.

228 Throughout tasting, consumers were told to drink the water provided to ensure palate
229 cleansing.

230 *2.6. Statistical Analysis*

231 *2.6.1 Chemical composition*

232 Data were analyzed using one-way Analysis of Variance (ANOVA) with sample as
233 fixed factor and protein, moisture, iron, vitamin A, thickness, diameter, spread ratio,
234 breaking force, a^* , b^* , L^* , as dependent variables. When a significant sample effect
235 was obtained, a Student Neuman-Keuls pair comparison test was applied ($\alpha = 5\%$).

236 *2.6.2. Sensory evaluation*

237 Principal Component Analysis (PCA) was performed on the consumer x biscuit global
238 liking scores to evaluate the agreement among consumers' judgments. Consumer
239 segmentation was then obtained by performing a Hierarchical Cluster Analysis (HCA)
240 of the coordinates of the consumers on the first two dimensions of the PCA. A
241 Euclidean distance and a Ward aggregation criteria were used for the HCA.

242 For each consumer cluster generated by the HCA, an ANOVA was performed on all
243 hedonic scales (global liking, appearance, odor, taste, texture) considering consumer
244 as random factor and samples as fixed factor. When a significant effect of sample
245 was obtained a Student Neuman-Keuls (SNK) pair comparison test was applied with
246 an alpha risk of 5%.

247 The open-ended question was analyzed by grouping terms with similar meaning.

248 This was done by two researchers. For each cluster, we computed the number of
249 consumers who used each term to describe what they liked and disliked in each
250 biscuit. Only terms used by more than two consumers were considered in the rest of

251 the analysis. A correspondence analysis was performed to visualize the relationship
252 between the samples and the descriptions.

253 **3. Results**

254 *3.1 Biscuits characterization*

255 *3.1.1 Proximate analysis*

256 Protein, iron, vitamin A content and moisture of raw MDL and MDL supplemented
257 biscuits are presented in Table 1. The analysis of the raw MDL revealed a content of
258 4.48 g of moisture, 20.7 g of protein, 11.77 mg of iron, 0.14 mg of vitamin A in 100 g
259 of powder. These results are within the range of the percentage of protein (18.2 -
260 29.8 g/100 g of MDL) reported in Witt's meta-analysis. However, the percentage of
261 iron content is lower than the values given by this meta-analysis (21.72 - 43.28 mg /
262 100 g of MDL). The vitamin A content in C1, C2 and C3 are below the detection limit.
263 This result suggests that vitamin A was degraded during baking.

264 The ANOVA result showed a significant effect of MDL concentration on moisture (F
265 (3; 8) = 129.99, $p < 0.0001$), protein (F (3; 8) = 79.57, $p < 0.0001$) and iron (F (3; 8) =
266 708.63, $p < 0.0001$) composition of biscuits. The pair comparison test indicates that
267 the control biscuit C0 is significantly different in moisture content from the biscuits
268 containing MDL. The amount of moisture increases with the concentration of MDL but
269 the pair comparison test shows no significant difference between C1 and C2. The
270 gradual increase of MDL concentration added to biscuits also caused a gradual
271 increase in protein content. The pair comparison test, revealed a significant
272 difference between the control C0 and all other biscuits. Thereby, a substitution of
273 wheat flour with MDL (from 10 to 30%) increases the protein content in biscuit by plus

274 10, 17, and 21% (for C1, C2 and C3 respectively) based on the control biscuit.
275 Finally, the addition of MDL increases significantly the biscuit iron content from 0.85
276 mg/100 g (biscuit C1) to 2.46 mg/100 g (biscuit C3). The pair comparison shows a
277 significant difference between all biscuits.

278 *3.1.2 In vitro protein digestibility (IVPD)*

279 The percentage of enzymatic hydrolysis (%EH) of casein is 92%, which corresponds
280 to the expected 100% of IVPD of casein and thus validates our protocol. For raw
281 MDL we obtained 57% of IVPD. The IVPD of biscuit C1 (84%) was higher than the
282 control biscuit C0 (47%). Then, the IVPD decreased from C1, to C2 (58%), to C3
283 (54%).

284 *3.1.3 Biscuit physical properties*

285 The analysis of the physical characteristics of the supplemented biscuits is presented
286 in Table 1. The ANOVA indicates that there is a significant difference between
287 biscuits for all parameters. A significant decrease in thickness was observed with the
288 increase in MDL concentration ($F(3; 8) = 162.88; p < 0.0001$). The biscuit containing
289 the highest concentration of MDL (C3) exhibited the lowest thickness of 6.72 ± 0.19
290 mm, whereas the control biscuit (C0) showed the highest thickness of 9.44 ± 0.10
291 mm.

292 The diameter of the C3 biscuit was significantly higher ($F(3; 8) = 35.45; p < 0.0001$)
293 than the diameter of the control sample while no significant difference was observed
294 among C0, C1 and C2 samples. The spread ratio also increased significantly as the
295 level of MDL added to the biscuit increased ($F(3; 8) = 113.86; p < 0.0001$).

296 The breaking force increased significantly with the concentration of MDL ($F(3; 8) =$
297 $160.05; p < 0.0001$). A weight of 15.44 kg was necessary to break the biscuit with the

298 highest MDL concentration (C3), whereas, only 3.08 kg was necessary to break the
299 biscuit without MDL. The pair comparison test showed no significant difference
300 between samples C1 and C2.

301 The impact of incorporation of MDL on biscuit color parameters is presented in Table
302 1. The ANOVA showed a significant effect of MDL on L* values ($F(3; 8) = 227.01$; $p < 0.0001$). The SNK test showed that the lightness parameter significantly decreased
303 (< 0.0001). The SNK test showed that the lightness parameter significantly decreased
304 (from 75.06 to 41.67) with increase in MDL concentration: The more the quantity of
305 MDL added to the control sample, the darker the biscuits. The a* value parameter
306 also reflects a significant effect ($F(3; 8) = 141.71$; $p < 0.0001$) on biscuits containing
307 MDL. The increase of MDL concentration moved biscuits from the red color area
308 (+a*) to the green color area (-a*). For C0, an a* value of +1.9 was observed, while
309 the a* values of C1, C2 and C3 were respectively -0.13; -0.27; -1.53. The pair
310 comparison test shows no difference between C1 and C2 in term of a*. The ANOVA
311 showed that the b* value decreased significantly ($F(3; 8) = 192.37$; $p < 0.0001$) from
312 C0 to C3 (21.43 to 7.25) indicating that with the addition of MDL, biscuits had a dark
313 blue color.

314 *3.2. Sensory evaluation results*

315 *3.2.1. Consumer segmentation*

316 The first two Principal Components (PC) of the PCA explain 74.9% of the total
317 variance of the overall liking scores (Fig. 2a and 2b). All four samples load positively
318 on the first PC (53.4%). Consumers that are positively correlated with this component
319 tend to like better the supplemented biscuits than other consumers. The second PC
320 (21.5%) opposes samples C0 and C3. Consumers that are positively correlated with

321 this PC preferred sample C3 to sample C0 and conversely, consumers negatively
322 correlated with this PC preferred sample C0 to sample C3.

323 The HCA separated the consumers into two clusters (Fig. 2b). Consumers from
324 cluster 1 (N=49) tend to like the biscuits less than consumers from cluster 2 (N=42).
325 The two clusters differ significantly in term of consumers' age ($F(1; 89) = 5.53$; p
326 < 0.0209). Consumers in cluster 2 are much younger (median age = 24.5) than the
327 consumers in cluster 1 (median age = 26). No significant difference was observed in
328 term of nationality ($\chi^2 = 1.33$) or gender ($\chi^2 = 0.26$).

329 Fig 3a and b represent the global liking score distributions for cluster 1 and 2.
330 Consumers from cluster 1 (Fig. 3a) accepted only the control biscuit C0 (median = 5).
331 Seventy-five percent of consumers rated the C1 biscuit between 2 and 4 with a
332 median of 4.3. For C2 and C3 products, all scores were below 4 with a median of 3
333 and 2 respectively. Consumers from cluster 2 (Fig. 3b) liked better the biscuits
334 containing MDL than consumers from cluster 1 (average score > 4). For the C0
335 product, 75% of consumers had liking scores greater than 5 with a median of 6. For
336 product C1 and C2, 50% of consumers had scores between 5 and 6 and 4 and 5 with
337 medians equal to 6 and 5 respectively. The liking scores for C3 were more variable
338 with some consumers clearly disliking this biscuit (scores close to 1) and other liking
339 it (scores equal to 6).

340 *3.2.2. Effect of Moringa content on overall, appearance, texture, smell and taste liking* 341 *scores*

342 ANOVA and pairwise comparison tests were carried out separately for each cluster
343 and each rating scale (Table 2). A sample effect was observed in both clusters for
344 each rating scale. This effect is, however, stronger in the first cluster than in the

345 second one. In both clusters, appearance is the factor that affects the liking of the
346 biscuit the most ($R^2 = 0.76$ and 0.71 for cluster 1 and 2 respectively). In the first
347 cluster, texture is the factor that has least effects on the liking of the biscuit. The other
348 scales have comparable effects. In the second cluster, texture influences the liking
349 scores more than taste and odor, which is the least impacting factor.

350 The pair comparison indicates that for both clusters and all liking scales the control
351 biscuit is significantly preferred to the MDL supplemented biscuits. In the first cluster,
352 C2 and C3 are not significantly different for global liking, odor, and texture scale. A
353 significant difference was observed for appearance and taste. In the second cluster,
354 we observe significant differences between all samples for all scales but odors.

355 *3.2.3. Analysis of open-ended question*

356 We represented the frequencies of positive and negative terms as word clouds
357 separately for each cluster (Fig. 4). For cluster 1, (consumers who liked less the
358 Moringa supplemented biscuits) the most frequent word used to describe what
359 consumers liked in the four samples is a texture term: crunchy. Consumers liked also
360 the taste, especially the sweetness of the control sample as well as its color. On the
361 negative side, they found it bland. Color was the most frequently used term to
362 describe what consumers disliked in the samples supplemented with MDL. The term
363 too hard was also mentioned for samples C2 and C3 followed by vegetal. For sample
364 C3, terms frequently mentioned include odor, spinach, unpleasant, too strong, and
365 bitter.

366 For Cluster 2, (consumers who liked more the MDL supplemented biscuits) crunchy
367 was frequently used to describe the positive aspects of the four biscuits as for cluster
368 1. Sweet and color were also mentioned frequently for C0 whereas new terms appear

369 to describe what consumers liked in the MDL supplemented biscuit such as taste,
370 sweet, color and crispy for sample C1. Cluster 2 consumers also appreciated the
371 originality of the MDL supplemented biscuits, especially in sample C3. Finally, the
372 term vegetal was mentioned among the positive aspects of sample C2. For the
373 negative side, as in cluster 1, consumers cited the term bland for sample C0. For the
374 other samples, color was much less cited than in cluster 1. For sample C1 the most
375 frequent term was vegetal. After-taste and spinach were also important for this
376 sample. C2 was characterized mostly by the term too hard, followed by color, vegetal
377 and bitter. C3 was perceived as having a taste that is too strong in addition to being
378 vegetal and bitter. Color was also cited but in a lesser extent than for samples C2
379 and C1.

380 Fig. 5 shows the first two dimensions (71.5% of the variance) of the CA carried out on
381 the term frequency counts. For both clusters, the first dimension (55.5%) opposes the
382 control sample to the MDL supplemented samples. In both clusters, the control
383 sample is positively perceived except for the term bland whereas the MDL
384 supplemented samples have negative global perception among consumers. The
385 second dimension opposes the two clusters. The difference between the clusters
386 increases with the proportion of MDL included in the biscuit. In cluster 1, the MDL
387 supplemented biscuits attracted mostly negative terms, whereas in cluster 2 some
388 positive terms were associated to these biscuits. Sample C1 attracted terms such as
389 pleasant and balanced and samples C2 and C3 were positively associated with terms
390 such as vegetal and original. Consumers in this group diverge somewhat in their
391 appreciation of sample C3. While some consumers liked its originality and its vegetal
392 character, others were less positive and associated it with negative surprise,
393 bitterness and find its taste too strong.

394 **4. Discussion**

395 The first objective of this work was to implement sensory analysis in order to gain
396 understanding of heterogeneity in consumer preferences of Moringa supplemented
397 biscuits. Secondly, this study also investigated nutritional enrichment of biscuits
398 supplemented with Moringa.

399 *4.1 Effects of MDL supplementation on biscuits nutritional properties and IVPD*

400 Although raw MDL contain vitamin A, this compound was degraded during the baking
401 process. In order to avoid this loss, one solution could be to incorporate MDL after
402 the cooking process (e.g. soup, sauce) or in raw products (e.g. salad). However, this
403 solution may lead to nutritional or sensory related problems. Findings from the
404 sensory analysis revealed that MDL had a pronounced vegetal taste, which is
405 unpleasant when eaten raw. Previous studies showed that heat processing was an
406 effective method to inactivate anti-nutritional factors of most tropical legumes. The
407 inactivation of trypsin inhibitor, for example, makes protein more susceptible to
408 enzymatic digestion (Akande and Fabiyi, 2010). Therefore, another solution could be
409 to use cooking processes with low heating impact in order to find a trade-off between
410 all these criteria.

411 Although, in agreement with previous work (Dachana et al., 2010; Mouminah, 2019;
412 Rathnayake and Navarathna, 2017), the addition of MDL increased protein and iron
413 content of biscuits, only a percentage of these proteins were digested. With our IVPD
414 protocol (gastric digestion 2h; intestinal digestion 2h), 57% of the proteins of raw
415 MDL was digested. This value is within the range of IVPD values reported by Teixeira
416 et al. (2013) (33.3%) and Ly et al. (2001) (74.2%). This difference in percentage of
417 IVPD can be explained by the difference in experimental conditions where, for

418 example, for Teixeira et al. (2013) the gastric digestion takes 3h and the intestinal
419 digestion takes 1h.

420 Moreover, with respect to the IVPD of biscuits, the results showed that the biscuit
421 sample which had the lowest concentration of MDL (C1), had the highest IVPD of
422 84%. Beyond this concentration, the IVPD decreases when the protein concentration
423 in biscuits increases. This decrease of digestibility could be explained by an
424 incomplete enzymatic hydrolysis of proteins due to insufficiency of digestion time or
425 enzyme concentration. Nevertheless, Boisen and Fernandez (1995) confirmed that in
426 vitro digestibility can predict in vivo ideal digestibility. Thus, the appropriate solution is
427 maybe to avoid incorporating a fairly large amount of MDL in the biscuit in order to
428 optimize their digestibility and their benefits.

429 *4.2. Contribution of supplemented biscuit to recommended daily requirement*

430 In order to determine the contribution of biscuits containing MDL to the recommended
431 daily requirement, we chose the C1 biscuit, which is the best digested and the most
432 widely accepted by consumers. So, 100 g of biscuit C1 (10%MDL; 84%IVPD) /day
433 provide about 0.85 mg of iron and 5.11 g of proteins. Compared with the
434 recommended daily requirement of adults (11 mg of iron /day and 0.83 g of
435 protein/kg/day (ANSES, 2016), the consumption of 100 g of biscuits C1/ day would
436 provide 7.7% of the iron. For protein, this consumption provides just 8.5% of the
437 protein after digestion (for an adult weighting 60 kg). Despite this rather low
438 percentage of protein intake, the leaves of Moringa contain the eight essential amino
439 acids among which the most abundant is leucine and the least abundant is
440 methionine (Ramaroson et al., 2014). Wheat protein have a lower nutritional quality
441 since they are deficient in essential amino acids such as lysine and threonine

442 (Dewettinck et al., 2008; Ndife et al., 2011). Combining MDL with wheat flour
443 products can improve essential amino acids composition.

444 *4.3. Effects of MDL supplementation on physical properties of biscuits*

445 The incorporation of MDL in biscuits impacts also the physical properties of biscuits,
446 as thickness decreases. This change is due to the reduction of the percentage of
447 gluten contained in the blend. It has been reported by Sengev et al. (2013) that
448 partial replacement of wheat flour with non-glutinous flour results in lower bread
449 volumes. Gluten is the main structure-forming protein in flour. It is responsible for the
450 elastic and extensible properties needed to produce good quality bread (Greene and
451 Bovell-Benjamin, 2004). In addition, the breaking force increases when the
452 percentage of MDL incorporated in the biscuits increases. This explains why the
453 texture of the supplemented biscuit was affected by the addition of MDL. This result
454 is in line with Galla et al. (2017) who show that the increase of leafy spinach in
455 biscuits increases hardness and breaking strength. All of these changes and
456 constraints confirm that there is a limit to further increasing the amount of MDL in the
457 biscuit in order to provide more nutrients.

458 *4.4. Effects of MDL supplementation on biscuit sensory properties and consumer* 459 *acceptance*

460 The results of this study show that the global liking of biscuits containing MDL was
461 not homogenous. Some consumers (cluster 2) tended to like biscuits containing MDL
462 whereas, other consumers did not accept them (cluster 1). When considering the
463 different sensory modalities, the appearance was the factor that affected the most the
464 global liking of biscuits for both consumer groups. The open-ended question
465 suggests that color is the main visual driver of rejecting biscuits supplemented with
466 MDL: color was clearly disliked by consumers of cluster 1 but less so by consumers

467 from cluster 2. Consumers in cluster 2 were younger than consumers in cluster 1.
468 This age difference perhaps implies that younger consumers are more willing to
469 accept the supplemented biscuit due to their curiosity to test novel products as
470 attested by the high frequency of the term original in this cluster. This interpretation is
471 in line with the study of Ramaroson et al. (2014) carried out in Madagascar.
472 Concerning cluster 1, color hampers consumer's acceptance of MDL supplemented
473 biscuit. This might be explained by the fact that consumers in this cluster were not
474 familiar with the green color brought by MDL. The rejection of MDL supplemented
475 biscuits by these consumers could thus be interpreted in terms of food neophobia,
476 which is the avoidance of, or reluctance to consume novel foods (Aldridge et al.,
477 2009). Texture was also one of the most important drivers of biscuit rejection. The
478 main problem was the hardness of the biscuit: the force needed to break the biscuit
479 sample that had the highest concentration of MDL (C3) was five times higher than the
480 force needed to break the control biscuit. In line with this interpretation, the term too
481 hard was frequently evoked in the open-ended question for MDL supplemented
482 biscuits.

483 **5. Conclusion**

484 The present study showed that the incorporation of MDL in food products like biscuits
485 would encounter several obstacles to consumer acceptance and affect physical
486 properties especially in terms of color (green) and texture (hardness). For nutritional
487 benefits, this fortification can enhance nutritional values such as iron and protein
488 compounds but many constraints and negative effects appear following this
489 incorporation. Protein provided by these biscuits were not completely digested.
490 Besides, due to baking processes, some of the nutrients from Moringa such as

491 vitamin A can be lost. Therefore, incorporating a high concentration of MDL in
492 biscuits is constrained by a lot of limits as previously explained.

493 In conclusion, *Moringa oleifera* could be a proper source to enhance nutritional
494 compounds in foods. In addition, some advice can be given to food producers who
495 are interested in food supplementation: it is essential to choose the appropriate
496 product in which to incorporate Moringa leaves to benefit from these dietary
497 advantages with the right quantity without influencing the technological and sensory
498 properties of products, mainly texture and appearance.

499 **Declaration of Competing Interest:** The authors declare no conflict of interest.

500 **Acknowledgements**

501 The authors would like to thank AgroSup Dijon and Embassy of Tunisia in France for
502 the financial support of this work. We gratefully thank Bernadette Rollin, Anne-Sophie
503 Viaux and Christine Rojas for their assistance with the laboratory work. Catalina
504 Onofrei is particularly acknowledged for English proofreading.

505 **References**

506 Abdel-Samie, M., Abdulla, G., 2014. Effect of *Moringa oleifera* leaves (*Moringa*
507 *oleifera* Lam.) on some physico-chemical and sensory properties of wheat flour
508 cookies. Zagazig Journal of Agricultural Research, 41, 304-315.

509
510 Adeyemi, S.B., Ogundele, K.O., Animasaun, M.A., 2014. Influence of drying methods
511 on the proximate and phytochemical composition of *Moringa oleifera* Lam. Global
512 Journal of Medicinal Plant Research, 2, 1-5.

513
514 Akande, K.E.; Fabiyi, E.F., 2010. Effect of processing methods on some
515 antinutritional factors in legume seeds for poultry feeding. International Journal of
516 Poultry Science, 9, 966-1001.

517
518 Aldridge, V., Dovey, T.M., Halford, J.C.G., 2009. The role of familiarity in dietary
519 development. Developmental Review, 29, 32-44.

520

521 Arshad, M.U., Faqir, M.A., Tahir, Z., 2007. Nutritional assessment of cookies
522 supplemented with defatted wheat germ. Food Chemistry, 102, 123-128.
523

524 ANSES, 2016. Agence nationale de sécurité sanitaire de l'alimentation, de
525 l'environnement et du travail. <https://www.anses.fr/fr/content/le-fer/> Accessed 2019,
526 November, 18.
527

528 Boisen, S., Fernandez, J.A., 1995. Prediction of the apparent ileal digestibility of
529 protein and amino acids in feedstuffs and feed mixtures for pigs by in vitro analyses.
530 Animal Feed Science and Technology, 51, 29-43.
531

532 Bruins, M., Bird, J., Aebischer, C., Eggersdorfer, M., 2018. Considerations for
533 secondary prevention of nutritional deficiencies in high-risk groups in high-income
534 countries. Nutrients, 10, 2-15.
535

536 CBI, 2019. Centre for the promotion of imports from developing countries, exporting
537 Moringa to Europe. [https://www.cbi.eu/market-information/natural-ingredients-health-
538 products/moringa/](https://www.cbi.eu/market-information/natural-ingredients-health-products/moringa/) Accessed 2019, September, 07.
539

540 Daba, M., 2016. Miracle tree: A review on multi-purposes of *Moringa oleifera* and its
541 implication for climate change mitigation. Journal of Earth Science & Climatic Change
542 7, 366.
543

544 Dachana, K.B., Jyotsna, R., Indrani, D., Prakash, J., 2010. Effect of dried Moringa
545 (*Moringa Oleifera* Lam.) leaves on rheological, microstructural, nutritional, textural
546 and organoleptic characteristics of cookies. Journal of Food Quality, 33, 660-677.
547

548 Dewettinck, K.F., Van Bockstaele, B., Kühne, D., Van de Walle, T., Courtens, M.,
549 Gellynck, X., 2008. Nutritional value of bread: Influence of processing, food
550 interaction and consumer perception. Journal of Cereal Science, 48, 243-257.
551

552 Fahey, J., 2005. *Moringa oleifera*: A review of the medical evidence for its nutritional,
553 therapeutic, and prophylactic properties. Trees for Life Journal.
554

555 Ganorkar, P.M., Jain, R.K., 2014. Effect of flaxseed incorporation on physical,
556 textural, sensorial and chemical attributes of cookies. International Food Research
557 Journal, 21, 1515-1521.
558

559 Galla, N.R., Pamidighantam, P.R., Karakala, B., Gurusiddaiah, M.G., Akula, S., 2017.
560 Nutritional, textural and sensory quality of biscuits supplemented with spinach
561 (*Spinacia oleracea* L.). International Journal of Gastronomy and Food Science, 7,
562 20-26.
563

564 Greene, J.L., Bovell-Benjamin, A.C., 2004. Macroscopic and sensory evaluation of
565 bread supplemented with sweet-potato flour. Journal of Food Science, 69, 167-173.
566

567
568 Hekmat, S., Kathryn, M., Soltani, M., Robert, G., 2015. Sensory evaluation of locally-
569 grown fruit purees and inulin fibre on probiotic yogurt in Mwanza, Tanzania and the
570 microbial analysis of probiotic yogurt fortified with *Moringa oleifera*. Journal of Health,
571 Population, and Nutrition, 33, 60-67.
572
573
574 Lee, J., Chambers, D.H., Chambers, E., 2014. A comparison of the flavor of green
575 teas from around the world. Journal of the Science of Food and Agriculture, 94,
576 1315-1324.
577
578
579 Ly, J., Samkol, P., Preston, T.R., 2001. Nutritional evaluation of tropical leaves for
580 pigs: Pepsin/pancreatin digestibility of thirteen plant species. Livestock Research for
581 Rural Development. <http://www.lrrd.org/lrrd13/4/ly134a.htm>
582
583
584 Minekus, M., Alming, M., Alvito, P., Balance, S., Bohn, T., Bourlieu, C., Carrière, F.,
585 2014. A standardised static in vitro digestion method suitable for food – an
586 international consensus. Food & Function, 5, 1113-1124.
587
588 Mouminah, S.H., 2019. Effect of dried *Moringa oleifera* leaves on the nutritional and
589 organoleptic characteristics of cookies. International Journal of Science and
590 Agricultural Environments, 36, 297-302.
591
592 Nagi, H.P.S., Kaur, J., Dar, B.N., Sharma, S., 2012. Effect of storage period and
593 packaging on the shelf life of cereal bran incorporated biscuits. American Journal of
594 Food Technology, 7, 301-310.
595
596 Ndife, J., Abdulraheem, L.O., Zakari, U.M., 2011. Evaluation of the nutritional and
597 sensory quality of functional breads produced from whole wheat and soya bean flour
598 blends. African Journal of Food Science, 5, 466-472.
599
600
601 Oyeyinka, A.T. Oyeyinka, A.S., 2016. *Moringa oleifera* as a food fortificant: Recent
602 trends and prospects. Journal of the Saudi Society of Agricultural Sciences, 17, 127-
603 136.
604
605 Ramarosan, R.V., Arvisenet, G., Valentin, D., 2014. Studying the nutritional beliefs
606 and food practices of Malagasy school children parents. A contribution to the
607 understanding of malnutrition in Madagascar. Appetite, 81, 67-75.
608
609 Ramota, O., Rowland, M.O., Oyeyinka, S.A., Adewumi, T., 2013. Proximate, mineral
610 and sensory qualities of 'Amala' prepared from Yam flour fortified with *Moringa* leaf
611 powder. Food Science and Quality Management, 12, 10-22.
612
613 Rathnayake, A.R.M.A.N., Navarathna, S.B., 2017. Utilization of *Moringa oleifera*
614 leaves as a functional food ingredient in bakery industry. International Journal of
615 Science and Research, 6, 339-344.

616
617 Salem, A.S., Salama, W., El Kebir, G., 2013. Enhancement of nutritional and
618 biological values of Labneh by adding dry leaves of *Moringa oleifera* as innovative
619 dairy products. World Applied Sciences Journal, 22, 1594-1602.

620 Sánchez-Machado, D. I., Núñez-Gastélum, J.A., Cuauhtémoc, R.M., Ramírez-Wong,
621 B., López-Cervantes, J., 2010. Nutritional quality of edible parts of *Moringa oleifera*.
622 Food Analytical Methods, 3, 175-180.

623
624 Siddhuraju, P., Becker, K., 2003. Antioxidant properties of various solvent extracts of
625 total phenolic constituents from three different agroclimatic origins of Drumstick tree
626 (*Moringa oleifera* Lam.) leaves. Journal of Agricultural and Food Chemistry, 51,
627 2144-2155.
628

629 Sengeev, A.I., Joseph, O.A., Gernah, D.I., 2013. Effect of *Moringa oleifera* leaf
630 powder supplementation on some quality characteristics of wheat bread. Food and
631 Nutrition Sciences, 4, 720-726.
632

633 Sreelatha, S., Padma, P.R., 2009. Antioxidant activity and total phenolic content of
634 *Moringa oleifera* leaves in two stages of maturity. Plant Foods for Human Nutrition,
635 64, 303-311.
636
637

638 Sulieman, A.A., Ke-Xue, Z., Wei, P., Hassan, H.A., Mohammed, O., Azhari, S., Hui-
639 Ming, Z., 2019. Rheological and quality characteristics of composite gluten-free
640 dough and biscuits supplemented with fermented and unfermented *Agaricus*
641 *bisporus* polysaccharide flour. Food Chemistry, 271, 193-203.
642
643

644 Symoneaux, R., Galmarini, M.V., 2014. Open-ended questions, in: Varela, P.,
645 Gastón, A. (Eds.) Novel Techniques in sensory characterization and consumer
646 profiling. Taylor & Francis group, London, pp. 307-330.
647
648

649 Teixeira, E.M., Carvalho, M.R., Neves, V.A., Silva, M.A., Arantes-Pereira, L., 2014.
650 Chemical characteristics and fractionation of proteins from *Moringa oleifera* Lam.
651 leaves. Food Chemistry, 147:51-54. <https://doi.org/10.1016/j.foodchem.2013.09.135>
652
653

654 Witt, K.A., 2013. The nutrient content of *Moringa oleifera* leaves, Ecocommunity.
655 <https://www.echocommunity.org/resources/a7ee06e3-40f2-4ef0-859e-4e64b90a56c8>
656
657

658 **Tables**

659

660 **Table 1** Physico-chemical characteristics of MDL powder and MDL supplemented
661 biscuits

	MDL	C0 (0% MDL)	C1 (10% MDL)	C2 (20% MDL)	C3 (30% MDL)
Moisture (g/100g)	4.48 ± 0.03	3.78 ± 0.11 ^b	2.71 ± 0.09 ^a	4.19 ± 0.05 ^c	4.64 ± 0.07 ^d
Proteins (g/100g)	20.70 ± 0.91	4.60 ± 0.09 ^a	5.11 ± 0.06 ^b	5.57 ± 0.20 ^c	5.96 ± 0.04 ^d
Iron (mg/100g)	11.77 ± 0.86	0.00 ^a	0.85 ± 0.07 ^b	1.72 ± 0.06 ^c	2.46 ± 0.10 ^d
Vitamin A (mg/100g)	0.14 ± 0.02	-	-	-	-
Thickness (mm)		9.44 ± 0.10 ^d	8.67 ± 0.17 ^c	7.39 ± 0.19 ^b	6.72 ± 0.19 ^a
Diameter (mm)		27.70 ± 0.10 ^a	27.47 ± 0.29 ^a	27.67 ± 0.15 ^a	28.83 ± 0.12 ^b
Spread ratio		29.33 ± 0.40 ^a	31.70 ± 0.92 ^b	37.46 ± 1.05 ^c	42.92 ± 1.34 ^d
Breaking force (Kg)		3.08 ± 0.40 ^a	7.78 ± 0.76 ^b	7.93 ± 0.41 ^b	15.44 ± 1.03 ^c
L*		75.06 ± 3.08 ^d	53.59 ± 1.29 ^c	44.92 ± 0.48 ^b	41.67 ± 0.74 ^a
a*		1.90 ± 0.03 ^c	-0.13 ± 0.01 ^b	-0.27 ± 0.01 ^b	-1.53 ± 0.41 ^a
b*		21.43 ± 1.17 ^d	16.78 ± 0.57 ^c	10.16 ± 0.88 ^b	7.25 ± 0.29 ^a

MDL: Moringa Dried Leaves. Values represent mean ± SD of triplicate assays. Means ± SD with the same letter are not significantly different at an alpha risk of 5%.

662

663 **Table 2** One-way repeated measure ANOVA with consumers as a random factor,
 664 samples as a fixed factor and liking scores as dependent variables.

Cluster 1	F	P	R²	C0	C1	C2	C3
Global liking	66.10	0.0001	0.65	4.9 ^a	3.7 ^b	2.6 ^c	2.4 ^c
Appearance	94.70	0.0001	0.76	5.6 ^a	3.9 ^b	3.0 ^c	2.3 ^d
Odor	45.69	0.0001	0.65	4.5 ^a	3.2 ^b	2.6 ^c	2.4 ^c
Texture	30.90	0.0001	0.60	5.3 ^a	4.7 ^b	3.4 ^c	3.4 ^c
Taste	69.20	0.0001	0.67	5.2 ^a	3.8 ^b	2.6 ^c	2.2 ^d
Cluster 2							
Global liking	24.60	0.0001	0.47	6.0 ^a	5.3 ^b	4.6 ^c	4.0 ^d
Appearance	54.97	0.0001	0.71	5.7 ^a	4.7 ^b	3.7 ^c	3.1 ^d
Odor	17.98	0.0001	0.48	5.2 ^a	4.0 ^b	3.7 ^{bc}	3.3 ^c
Texture	30.70	0.0001	0.61	6.0 ^a	5.4 ^b	4.1 ^c	3.9 ^c
Taste	27.39	0.0001	0.56	5.9 ^a	5.1 ^b	4.3 ^c	3.7 ^d

Means with the same letter are not significantly different at an alpha risk of 5%.

665 **Figure Captions**

666


667 **Fig. 1** Photographs of control biscuit (C0) and Moringa supplemented biscuits (C1,
668 C2 and C3)

669 **Fig. 2** Principal Components Analysis (PCA) of the consumers global liking scores.
670 The left panel (a) represents the loading of the samples and the right panel (b) the
671 projection of the consumers. The dot colors and the ellipses represent the two
672 clusters yielded by the Hierarchical Cluster Analysis (HCA)

673 **Fig. 3** Boxplots representing the frequency counts of global liking scores for
674 consumers from cluster 1(a) and cluster 2(b)

675 **Fig. 4** Word clouds of the terms generated by the consumers from the two clusters to
676 describe what they liked and disliked in the samples. The size of the words in this
677 representation is proportional to their use: the larger the word size, the higher the
678 frequency of occurrence

679 **Fig. 5** First 2-dimensional space of the correspondence analysis of the terms
680 generated by the consumers to describe what they like and dislike in the biscuits. The
681 size of the diamonds is proportional to the quality of the representation of the terms in
682 the subspace. Terms corresponding to what consumers like are followed by a sign
683 (+) and terms corresponding to what consumers disliked are followed by a sign (-).
684 Only terms used by at least 10% of consumers have been kept. Sample descriptions
685 obtained for consumers from cluster 1 are denoted by CL1 before the code of the
686 samples and sample descriptions obtained for consumers from cluster 2 are denoted
687 by CL2 before the code of the samples





C0


C1


C2

C3









Cluster1


C0

C1





C2

C3

Like



Dislike



Cluster2



C0

C1

C2

C3

Like



Dislike

