



HAL
open science

Potential of Meta-Omics to Provide Modern Microbial Indicators for Monitoring Soil Quality and Securing Food Production

Christophe Djemiel, Samuel S. Dequiedt, Battle Karimi, Aurélien Cottin, Walid Horrigue, Arthur Bailly, Ali Boutaleb, Sophie Sadet-Bourgeteau, Pierre-alain Maron, Nicolas Chemidlin Prévost-Bouré, et al.

► To cite this version:

Christophe Djemiel, Samuel S. Dequiedt, Battle Karimi, Aurélien Cottin, Walid Horrigue, et al.. Potential of Meta-Omics to Provide Modern Microbial Indicators for Monitoring Soil Quality and Securing Food Production. *Frontiers in Microbiology*, 2022, 13, pp.1-20. 10.3389/fmicb.2022.889788 . hal-03715998

HAL Id: hal-03715998

<https://institut-agro-dijon.hal.science/hal-03715998>

Submitted on 7 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OPEN ACCESS

Edited by:

Caroline Barretto,
Nestlé, Switzerland

Reviewed by:

Anna Gałazka,
Institute of Soil Science and Plant
Cultivation, Poland
Salma Mukhtar,
Connecticut Agricultural Experiment
Station, United States
Sandi Orlic,
Rudjer Boskovic Institute, Croatia

***Correspondence:**

Sébastien Terrat
sebastien.terrat@inrae.fr
Lionel Ranjard
lionel.ranjard@inrae.fr

†ORCID:

Christophe Djemiel
orcid.org/0000-0002-5659-7876
Battle Karimi
orcid.org/0000-0002-4523-974X
Aurélien Cottin
orcid.org/0000-0002-0862-784X
Walid Horrigue
orcid.org/0000-0001-9983-1039
Sophie Sadet-Bourgeteau
orcid.org/0000-0002-7533-4289
Pierre-Alain Maron
orcid.org/0000-0003-2315-0741
Nicolas Chemidlin Prévost-Bouré
orcid.org/0000-0002-8575-4295
Lionel Ranjard
orcid.org/0000-0002-7720-5843
Sébastien Terrat
orcid.org/0000-0001-5209-6196

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 04 March 2022

Accepted: 06 June 2022

Published: 30 June 2022

Citation:

Djemiel C, Dequiedt S, Karimi B,
Cottin A, Horrigue W, Bailly A,
Boutaleb A, Sadet-Bourgeteau S,
Maron P-A, Chemidlin
Prévost-Bouré N, Ranjard L and
Terrat S (2022) Potential of Meta-
Omics to Provide Modern Microbial
Indicators for Monitoring Soil Quality
and Securing Food Production.
Front. Microbiol. 13:889788.
doi: 10.3389/fmicb.2022.889788

Potential of Meta-Omics to Provide Modern Microbial Indicators for Monitoring Soil Quality and Securing Food Production

Christophe Djemiel^{1†}, Samuel Dequiedt¹, Battle Karimi^{1,2†}, Aurélien Cottin^{1†}, Walid Horrigue^{1†}, Arthur Bailly¹, Ali Boutaleb¹, Sophie Sadet-Bourgeteau^{1†}, Pierre-Alain Maron^{1†}, Nicolas Chemidlin Prévost-Bouré^{1†}, Lionel Ranjard^{1†*} and Sébastien Terrat^{1†*}

¹Agroécologie, INRAE, Institut Agro, Université Bourgogne, Université Bourgogne Franche-Comté, Dijon, France, ²Novasol Experts, Dijon, France

Soils are fundamental resources for agricultural production and play an essential role in food security. They represent the keystone of the food value chain because they harbor a large fraction of biodiversity—the backbone of the regulation of ecosystem services and “soil health” maintenance. In the face of the numerous causes of soil degradation such as unsustainable soil management practices, pollution, waste disposal, or the increasing number of extreme weather events, it has become clear that (i) preserving the soil biodiversity is key to food security, and (ii) biodiversity-based solutions for environmental monitoring have to be developed. Within the soil biodiversity reservoir, microbial diversity including Archaea, Bacteria, Fungi and protists is essential for ecosystem functioning and resilience. Microbial communities are also sensitive to various environmental drivers and to management practices; as a result, they are ideal candidates for monitoring soil quality assessment. The emergence of meta-omics approaches based on recent advances in high-throughput sequencing and bioinformatics has remarkably improved our ability to characterize microbial diversity and its potential functions. This revolution has substantially filled the knowledge gap about soil microbial diversity regulation and ecology, but also provided new and robust indicators of agricultural soil quality. We reviewed how meta-omics approaches replaced traditional methods and allowed developing modern microbial indicators of the soil biological quality. Each meta-omics approach is described in its general principles, methodologies, specificities, strengths and drawbacks, and illustrated with concrete applications for soil monitoring. The development of metabarcoding approaches in the last 20 years has led to a collection of microbial indicators that are now operational and available for the farming sector. Our review shows that despite the recent huge advances, some meta-omics approaches (e.g., metatranscriptomics or meta-proteomics) still need developments to be operational for environmental bio-monitoring. As regards prospects, we outline the importance of building up repositories of soil quality indicators. These are essential for objective and robust diagnosis, to help actors and stakeholders improve soil management,

with a view to or to contribute to combining the food and environmental quality of next-generation farming systems in the context of the agroecological transition.

Keywords: meta-omics, microbial indicators, food value chain, biomonitoring, soil quality

LINK BETWEEN SOIL AS A SUPPORT FOR FOOD PRODUCTION AND A HABITAT FOR LIVING ORGANISMS

Soil is the most complex natural physicochemical matrix and the most structured one on our planet. This heterogeneous layer is the most superficial one of the Earth's crusts, and can be from a few centimeters to several tens of meters deep. The variability of this matrix can be structured into three dimensions: horizontal, vertical, and temporal. Horizontal variability results from the spatial distribution of the different types of soils, influenced by the type of parent material (limestone, granite, metamorphic rock) on a large scale (Chesworth, 2007). Vertical variability refers to the different layers – called “horizons” – that have differentiated over time (Hartemink et al., 2020). Temporal variability results from the pedogenesis process, which is at the origin of soil creation from the weathering of the parent material by climate, vegetation and living organisms or human interventions. Soil formation is a slow process: 0.05 mm of soil are formed each year on average (Evans et al., 2019). Altogether, this makes soils a mosaic of extremely rich and diversified habitats providing a large number of macro- and micro-environments for micro-, meso- and macroscopic living organisms (Nielsen et al., 2015). It is home to many organisms that spend part or all of their life cycle underground and are part of the soil biota (e.g., nematodes, tardigrades, earthworms, microorganisms; **Figure 1**; Bardgett and Van Der Putten, 2014). Thanks to this mosaic of micro- and macro-habitats, soils host almost one third of our planet's biodiversity (FAOSTAT, 2021).

The food value chain summarizes the expression “from farm to fork,” including all steps from food production to food distribution to consumers, *via* food manufacturing. Soils are fundamental resources for agricultural production and play an essential role for food security (Kopittke et al., 2022). As providers of more than 95% of food and feed, they represent the cornerstone of the food value chain (Friedrichsen et al., 2019; FAOSTAT, 2021). Soil nutrient availability is unevenly distributed on regional and global scales and affects agricultural productivity (Silver et al., 2021). Soils harbor a large fraction of biodiversity – the backbone of the regulation of ecosystem services and soil multi-functionality. However, these are increasingly threatened by anthropogenic activities such as unsustainable soil management practices, pollution, waste disposal, or the increasing number of extreme weather events. Thus, monitoring the soil biodiversity has become an imperative and urgent issue to secure food production, and requires the development of microbial indicators (Kopittke et al., 2019). New initiatives have recently been launched at a global scale to try and build indicators of food system

sustainability (Béné et al., 2019), but microbial diversity was missing from the 27 selected indicators.

Soil Microorganisms: Modern Bioindicators of Soil Quality in a Context of Agroecological Transition


Definition of Bioindicators

A bioindicator (or biological indicator) is a measurement of macro- or micro-organisms whose response in terms of presence/absence, abundance, activities/functions, morphology, physiology or behavior provides information on the state of a habitat or ecosystem (Martinez-Salgado et al., 2010; Astudillo-García et al., 2019). It is used to assess the quality of an environment and its evolution over time. Bio-monitoring of soil quality is carried out at different spatial (local, landscape, regional or global) and temporal (years, decades, millennia) scales, but also at different levels of biological organization (individual, population, community, ecosystem; Pulleman et al., 2012).

Soil bioindicators are used to monitor soil responses to disturbances such as land use changes, agricultural management, or soil contamination, and to assess environmental risks (Geisen et al., 2019). It is essential for a bioindicator to have a good sensitivity to soil management and environmental pressures, and to be relevant to soil functions (e.g., organic matter decomposition, mineralization of soil nitrogen, formation of the soil structure) or related to soil quality (Bloem et al., 2005). From an operational perspective, an indicator should (i) be user-friendly in the field (ii) be precise and reproducible, and (iii) have an affordable financial cost to be used by the greatest number of actors and on a large scale (Griffiths et al., 2018). It should be as universal as possible for the various microbiomes (Schloter et al., 2018). Macro-fauna organisms (e.g., nematodes and earthworms) long remained the only indicators of soil biological quality because they were easily observed and well established in ISO standards (Martinez-Salgado et al., 2010; Schloter et al., 2018; Phillips et al., 2021). Microorganisms were used as bioindicators of soil quality later, but only when the molecular ecology era began (Maron et al., 2011). There are currently several bioindicators of soil quality available, based on microbial taxa and their associated functions (Schloter et al., 2018; Thiele-Bruhn et al., 2020).

Microorganisms Are Good Bioindicators of Soil Quality

Microorganisms (mainly bacteria, fungi and protists) and their functional guilds (pathotrophs, saprotrophs, symbiotrophs) are involved in many ecosystem services (**Figure 1**), in particular through their role in the biogeochemical cycles of major elements (Hermans et al., 2017; Frac et al., 2018; Geisen et al., 2018). They are involved in N cycling by fixing atmospheric N,




denitrification and ammonification (Hayatsu et al., 2008). Soil organic matter mineralization is mainly driven by microbial communities that transform the complex carbohydrates into mineral elements available for plant growth (Nielsen et al., 2015; Maron et al., 2018; Yang et al., 2018). Various microorganisms – especially bacteria and fungi – can achieve symbiosis with plants and benefit to plant growth and productivity (Prudent et al., 2020; **Figure 1**). They can also control plant pathogens thanks to metabolites and provide a barrier against pathogens (**Figure 1**). Soil microbial communities deliver other kinds of services that promote the soil structure or reduce soil contamination (FAO, ITPS, GSBI, and SCBD, 2020; **Figure 1**). Their deep involvement in ecological services make microorganisms ideal candidate bioindicators for monitoring the soil quality and food safety (**Figure 2**). In addition, their high sensitivity to environmental disturbances combined with their short generation time makes them early indicators of disturbances. As the cost of some meta-omics technologies is getting lower, the trends are now to define

new bioindicators, and an inventory of operational bioindicators is necessary to provide a benchmark for soil quality assessment. This evaluation needs to include (i) the currently available meta-omics approaches so as to evaluate their benefits and technical or conceptual drawbacks (ii) the establishment of an adapted sampling design, and (iii) the development of repositories for an objective interpretation of the results.

CONTRIBUTION OF -OMICS TECHNOLOGIES TO THE MONITORING OF THE SOIL MICROBIOLOGICAL QUALITY

Over the past two decades, many molecular tools have emerged that have accelerated the exploration of the soil microbial diversity, and later contributed to soil quality assessment



(Schloter et al., 2018; Nkongolo and Narendrula-Kotha, 2020). These methods can be divided into three major categories: (i) quantitative methods (ii) hybridization methods (microarrays), and (iii) sequencing methods.

Quantitative Methods

Different quantitative methods have been developed to investigate the relative or absolute abundance of taxa but also their functional profiles by targeting specific genes or gene families of interest (e.g., antibiotic resistome, functional genes in the N cycle; Oshiki et al., 2018). Quantitative polymerase chain reaction (qPCR), high-throughput quantitative PCR (HT-qPCR) and droplet digital PCR (ddPCR) provide access to the numbers of copies of one or more genes. Technological advances have made it possible to simultaneously analyze several thousands of samples and targets (Porter and Hajibabaei, 2018; Mehle and Dreo, 2019). More sensitive and more accurate, ddPCR appears to be promising for an absolute quantification of genes from complex samples such as soil samples because it can overcome the limitations associated with the presence of inhibitors (Gutiérrez-Aguirre et al., 2015).

Microarray Methods

DNA microarray is based on hybridization between (i) specific oligonucleotides used as probes and fixed on a solid surface, and (ii) the target corresponding to the DNA/RNA soil sample (Srivastava et al., 2019). Combining hybridization and labeling (e.g., by fluorescence) of DNA, this semi-quantitative technology is used to detect the presence or relative abundance of target markers or to monitor gene expression profiles at the individual or community levels (Sessitsch et al., 2006; Porter and Hajibabaei, 2018). Various microarrays such as PhyloChip, GeoChip (He et al., 2007), PathoChip (Lee et al., 2013), StressChip (Zhou et al., 2013), CAZyChip (Abot et al., 2016) have been developed to capture a snapshot of specific microbial populations and/or functional genes. For example, GeoChip was designed to detect genes from various metabolic pathways including the carbon, nitrogen, sulfur and phosphorus cycles. The latest version includes about 160,000 distinct probes covering about 1,500 functional gene families. PathoChip is less widespread and contains over 3,000 probes characterizing virulence factors (e.g., adhesion, colonization, motility, toxins) covering 1,400 species. CAZyChip was developed in 2016 to monitor the expression profiles of genes encoding bacterial enzymes specifically involved in polysaccharide degradation. These DNA

chips contain 55,220 probes targeting glycoside hydrolases covering more than 70% of the families present in the CAZy database. Some of these microarrays have become tools for diagnosis, in particular for targeting viruses, bacterial or fungal pathogens or harmful organisms.

From DNA Fingerprinting to High-Throughput Sequencing: Characterizing Microbial Diversity and Functions

DNA fingerprinting was the first high-throughput analytical technique used to comprehensively characterize the genetic structure of soil microbial communities in terms of presence, absence or relative abundance. These approaches are frequently based on PCR amplification of a molecular marker, and classified in two groups depending on the polymorphism of the studied sequences. The first group is usually based on the sequence length polymorphism and includes T-RFLP (restriction fragment length polymorphism), ARDRA [amplified rDNA (ribosomal DNA) restriction analysis], or ARISA (automated ribosomal intergenic spacer analysis). The second group relies on the polymorphism in the composition of the nucleotide sequence and involves techniques such as DGGE (denaturing gradient gel electrophoresis) or TGGE (temperature gradient gel electrophoresis; Rincon-Florez et al., 2013). These tools have been used for various soil microbial diagnostics because they are inexpensive and relatively fast (Sessitsch et al., 2006; Rincon-Florez et al., 2013; van Dorst et al., 2014). However, their sensitivity only identifies the dominant groups, and the comparison of samples (inter-gel) remains difficult (Schloter et al., 2018). Moreover, they do not provide any identification of the organisms or measurement of microbial diversity. These fingerprinting tools have been totally replaced by high-throughput sequencing.

Second- and third-generation high-throughput sequencing offers new ways of answering those various technical and scientific issues. Sequencing coverage, raw data quality, read length, technical biases vary depending on the method (Goodwin et al., 2016). Nevertheless, the performance of high-throughput sequencing platforms is constantly evolving to better support scientific goals at increasingly affordable costs. This has enabled the democratization of several meta-omics approaches for scientific teams but also the transfer of bio-indicators for the agri-food industries. In the next section, we explore and describe the different promising meta-omics approaches for the development of soil quality bioindicators (Figure 3).

META-OMICS APPROACHES: GUIDELINES FOR AND APPLICATIONS TO THE FOOD VALUE CHAIN

A paradigm shift emerged in environmental microbiology studies at the end of the 1990's, as the studies based on culture-dependent methods disappeared and were replaced by culture-independent methods. The transition was facilitated

by our capacity to extract and access the molecular resources (DNA, RNA, proteins, metabolites) of microbial communities directly from environmental samples. This determined the beginning of the present meta-omics area. These high-throughput approaches for screening hundreds to thousands of samples simultaneously and retrieving a huge amount of data have been developed and have evolved considerably over time. Studies related to soil and/or food have known rapid and important progress thanks to meta-omics since 2010 (Figure 4). However, very few studies related to soil and food were based on meta-omics compared to the number of studies focused on soil or food independently of each other. The metagenomic and metabarcoding surveys were the most frequently used approaches to study the soil and food microbiota and the microbiome, while metatranscriptomics and meta-proteomics were rarely investigated (Figure 4). The countries that produced the greatest numbers of publications about soil and/or food microbial research based on meta-omics were the USA and China, followed by Germany and France for European countries (Figure 5).

Assessing Analytical Specificity in Soil Analysis


Consequences of the Sampling Effort and Sample Conservation on Soil Analysis

Any sampling survey first involves a series of decisions that determine how the data should be analyzed (Dickie et al., 2018), e.g., what is sampled, what is not sampled, where it is sampled, how many samples to be taken, how many replicates, location of subsamples within plots, etc. Even storage conditions of soil samples can influence the quality of the molecular resource and should be well adapted. For example, the use of sieved soils and -40°C storage is efficient to access and keep DNA, but not to preserve and extract a more sensitive molecule like RNA (Ranjard et al., 2009). Fresh soils are undoubtedly an ideal material for characterizing microbial taxonomic and functional diversity, but using them is sometimes impossible in the case of large sampling surveys or comparisons among chronological samples requiring storage conditions (Cui et al., 2014; Table 1).

Extraction of Soil Genetic Resources: DNA and RNA

Meta-omics approaches started with isolating DNA directly from environmental samples. Success essentially depended upon the quality and the quantity of isolated DNA (Terrat et al., 2015). Various homemade DNA extraction protocols and commercial kits were developed to improve the efficiency of soil DNA extraction (Martin-Laurent et al., 2001; Delmont et al., 2011; Terrat et al., 2012). However, each method is characterized by its own advantages and biases, leading to variations in DNA representativeness. These variations have consequences on the observed effects on soil microbial assessments, and this makes it difficult to compare studies.

The choice of a specific DNA extraction method determines the inherent bias involved throughout the whole project




(Vestergaard et al., 2017). This is why standardized methods (like the ISO 11063: 2012 standard) are advised in order to provide consistent results across studies, more particularly when measurements are interpreted relatively to references used as bioindicators of soil quality (Table 1).

DNA provides information only about the presence of microorganisms in soils but does not inform on microbial activities. On the contrary, RNA analyses potentially describe what those microorganisms are actually doing and how fast they are doing it (Prosser, 2015). But RNA is a very sensitive molecule, with a low stability and a short half-life time (Hambraeus et al., 2003), and needs to be protected immediately after soil sampling using dedicated methods or solutions (e.g., LifeGuard Soil Preservation Solution or RNAlater). Moreover, to protect RNA from degradation by RNases, a clean laboratory must be dedicated. Inactivation reagents for RNases such as guanidine thiocyanate must be added to extraction buffers to inactivate RNase molecules immediately after their release from lysed cells. Finally, no method for RNA extraction from all types of soils is available;

this hinders the study of microbial gene expression in soils (Wang et al., 2012).

Extraction of Soil Molecular Resources: Proteins and Metabolites

Due to its huge potential for linking functional and phylogenetic information on soil microbial communities, research in soil ecology has showed a rising interest in meta-proteomics to investigate microbe-driven ecosystem functions (Prosser, 2015; Keiblinger et al., 2016). However, the application of meta-proteomics to soils still faces several challenges such as the heterogeneity of soil matrices or the ecosystem-specific dominance of a few microbial species, which limit the development of a generic protocol. The extraction of soil proteins is sensitive to the presence of several organic compounds such as complex carbohydrates, lipids, phenolics (e.g., lignin), humic substances, but also inorganic compounds such as silt and clay minerals (Keiblinger et al., 2016). Proteins may be adsorbed or linked anchored or embedded onto solid particles and soil



organic matter, hence reduced extraction efficiency. In addition, measurements of soil enzyme activities can be affected in different ways depending on the storage method, the soil type and even on the assay method. Freezing of small sampled aliquots retains the activity of most enzymes, but ultra-freezing is preferred for storage of samples with a high organic matter content (Wallenius et al., 2010). Altogether, processing fresh samples whenever possible or storage at -80°C is recommended to minimize the activity of naturally occurring proteases in environmental samples (Keiblinger et al., 2016).

Soil organic matter – more precisely the dissolved fraction – is composed of many small molecules of plant and microbial origin (Swenson et al., 2015). Classical methods for extracting soil metabolites involve long extraction times followed by compound-specific analyses (Swenson et al., 2015). However, the soil physical properties such as mineral composition, surface area, shape and porosity represent different challenges when it comes to extracting soil metabolites and metabolomics approaches. For example, amines adsorb to clay material in soils, and so do other metabolites, especially carboxylic acids that potentially form covalent bonds with organic matter or

are chelated by metals or other ions (from the soil or from the extractant). Moreover, each metabolite has specific half-life times depending on the soil specificities, but also on the activities of the organisms (i.e., enzymatic degradation). Finally, the quantity, stability and specificity of metabolites after extraction can be influenced by soil storage conditions (Table 1).

Metabarcoding to Investigate Microbial Diversity

Metabarcoding, also known as targeted metagenomics, uses the polymerase chain reaction (PCR) method to target a genomic region common to all studied microorganisms (Semenov, 2021). This molecular marker can be a gene allowing to study the relationships between microorganisms (considered as a taxonomic marker; Woese et al., 1990) or a functional gene/family (considered as a functional marker; Barbi et al., 2014). DNA fragments (amplicons) produced by the PCR and sequenced by high-throughput sequencing provide an overview of microbial diversity (Nilsson et al., 2019a). Several molecular markers exist to characterize




FIGURE 5 | Distribution of publications on meta-omics approaches in relation to soil (A), food (B) and soil and food (C). World distribution of publications on soil (A), food (B) and soil and food (C) using meta-omics approaches. This analysis relies on keywords present in the title or abstract, related to meta-omics (merging studies on metabarcoding, metagenomics, metatranscriptomics and meta-proteomics) and based on the country of the first author. Gray areas, absence of publication.

different microbial communities. The most popular marker is the gene encoding the small subunit ribosomal RNA (SSU rRNA), essential to ribosome formation. The 16S rRNA gene is used to determine the diversity of Archaea and Bacteria (Woese and Fox, 1977). Similar to the 16S rRNA gene, the 18S rRNA gene is used to explore eukaryotic organisms, and in particular the diversity of the fungal kingdom (Hadziavdic et al., 2014). Considered as the universal DNA barcode marker to investigate fungal diversity, the nuclear ribosomal internal transcribed spacer (ITS) is also applied

TABLE 1 | Influence of the conservation strategy of the soil sample on the quality of the extracted biological material.

Preservation strategy	DNA	RNA	Proteins	Metabolites
Immediate sample processing	+++	+++	+++	+++
Drying (room temperature)	+/-	-	-	-
Cold storage (4°C)	+/-	-	-	-
Freezing (-20°C)	++	+/-	+/-	+
Low freezing (-40°C to -80°C)	+++	+++	+++	+++
Chemical preservation (use of stabilizing solution)	+/-	++	++	unknown

+++, Image close to that of the sample; ++, Good protection of the sample; +, Average protection of the sample; +/-, Moderate impact of the storage method on the image; -, Strong impact of the storage method on the results.

to identify fungi. Among further taxonomic markers of other important functions are the *gyrB* (DNA gyrase subunit B), *recA* (recombinase A), or *rpoB* (RNA polymerase subunit beta) genes (Holmes et al., 2004; Poirier et al., 2018; Ogier et al., 2019). The choice of the marker depends on the targeted scientific question/objective and microbial communities. Using the metabarcoding approach is advantageous in several ways. It has become very popular in the last decade, and is undoubtedly the most affordable meta-omics approach. Thus, many projects have emerged to evaluate soil quality at large and local scales. Data analysis is now fully operational thanks to a significant range of open-source bioinformatics pipelines such as mothur (Schloss et al., 2009), qiime2 (Bolyen et al., 2019), FROGS (Escudie et al., 2018), or BIOCUM-PIPE (Djemiel et al., 2020a) and the associated standard operating procedures (SOPs) or tutorials. Nevertheless, this wide choice requires grasping well the subtleties of the algorithms implemented in the pipelines and their parameters because the accuracy of results is highly dependent on them (Pauvert et al., 2019). Another major effort in this area is the availability of reference databases for taxonomic assignment whatever the molecular marker (Quast et al., 2013; Nilsson et al., 2019b; Djemiel et al., 2020b). It should also be emphasized that some sequences could contain misidentifications or errors (Hofstetter et al., 2019). Therefore, it is important to be careful when interpreting taxonomic results, especially at the genus or species ranks. The initiation of a metabarcoding project requires choosing a molecular marker and then one or more regions to be amplified. This choice is crucial because the regions do not all have the same resolution and depend on the targeted samples and communities (Banos et al., 2018; Bukin et al., 2019). As a result, amplicon length is also crucial and impacts on the sequencers to be chosen and library preparation (Nilsson et al., 2019a; Castaño et al., 2020). The sequence length generally ranges from a few dozen to one hundred base pairs; this size does not allow fine taxonomic resolution to determine the species names. The use of metabarcoding approaches to monitor soil quality and also other compartments of the food chain value is scarce. However, some studies have already applied such approaches and brought significant knowledge on microbiota interactions

(Stefanini and Cavalieri, 2018). In a recent study, Zarraonaindia and colleagues investigated 725 samples from soils and grapevine organs using a metabarcoding approach to determine the influence of the vine cultivar, edaphic parameters, the vine developmental stage (dormancy, flowering, pre-harvest), and vineyard characteristics on the spatial and temporal dynamics of bacterial communities (Zarraonaindia et al., 2015). They concluded that most organ-associated taxa originated from the soil, and their distribution reflected the influence of highly localized biogeographic factors and vineyard management. Moreover, the vineyard-associated localization of bacterial taxa has implications for wine growers who rely on the assumption that the soil imparts a unique quality to the wine specific to each vineyard, called the “terroir effect.” Complementary to the evaluation of a microbial terroir, such studies investigate the environmental and human-related factors that influence the composition of microbial populations and potentially affect their performances, like fermentation (Stefanini and Cavalieri, 2018).

Metagenomics to Explore the Functional Potential of Microbial Communities

The metagenomics approach – or shotgun sequencing (without any *a priori* compared to metabarcoding) – rests upon the sequencing of total DNA from environmental samples. The extracted DNA is directly sequenced (or after fragmentation or nebulization). This avoids PCR amplification, and non-culturable and/or unknown organisms can be characterized, such as viruses (Prosser, 2015). Its main purpose is to characterize whole microbial genomes across microbiome samples. With the increasing efficiency of sequencing technologies, this approach can now be applied with a sufficient amount of acquired sequences to reconstruct “population genomes” of major microorganisms (Prosser, 2015). Metagenomics leads to an overview of the potential genomic structure of the microbial community, its potential functional properties, in a more sensitive manner than metabarcoding does (Bünemann et al., 2018). Moreover, with the democratization of sequencing technologies, a variety of tools and analysis pipelines have been developed to analyze metagenomics data (Kieser et al., 2020). Configuring various tools, linking them with advanced binning and annotation tools, and managing the updates and parameters modifications of each tool is now easier and more accessible for researchers with the efficient and automated deployment of workflows like ATLAS, MG-RAST or MOCAT2 (Meyer et al., 2008; Kultima et al., 2016; Kieser et al., 2020).

The use of metagenomics for characterizing microbial communities presents several advantages. As already stated, metagenomics characterizes unknown microorganisms in the absence of prior knowledge, and provides information on potential functions based on the genomic content. This allows a deeper understanding of the response of microbial communities to environmental changes (Aguilar-Pulido et al., 2016a). Some drawbacks have also been identified. The presence of a functional gene is not proof of its expression. The host organism may be dormant, inactive or only active when alternatively expressed

metabolic pathways do not require the function of the detected gene (Prosser, 2015). Moreover, for economic reasons the resulting datasets often have low coverage of the genomes in the microbiome. Finally, even with available bioinformatics pipelines, the analysis of such datasets needs expertise and can be time-consuming (Aguilar-Pulido et al., 2016a), with a high percentage of reads considered as “Unknown” due to the absence of similar reference genomes.

Metagenomics outcompetes metabarcoding in that it provides insights into the metabolic, virulence or resistance potential of microbial communities. A good example is the study of Wind and colleagues (Wind et al., 2021) focused on antibiotic resistance genes (ARG) across a vegetable production system. The study was interested on three pre-harvest “control-points,” i.e., manure and compost amendments, soils, and lettuce samples, to track antibiotic resistance and the associated microbiomes in order to identify the potential impacts of key agricultural management strategies. The soils were found highly stable in the composition of the resistome, which has the potential to act as a natural ecological buffer to ARG proliferation, at least over one harvest cycle. But lettuce grown in soils amended with compost produced from the manure of cows had higher total relative ARG abundance and risk scores compared to the soil amended with stockpiled manure. Similar results were described with poultry manure application: ARG abundance increased on lettuce surfaces at the greenhouse scale (Zhang et al., 2019). The authors of these studies advise reducing the spreading of ARG from amendment onto the lettuce leaf surface as this may influence human health negatively. Many other scientific questions can be tackled using metagenomics, such as Zn availability in grains for biofortification (Wang et al., 2021), or the detection of foodborne pathogens in production chains (Yang et al., 2016).

Metatranscriptomics Dedicated to Studying Gene Expression in Microbial Communities

Metatranscriptomics is quite similar to metagenomics, but different in that it focuses on microbial gene expression from complex environments. Metatranscriptomics provides the diversity of the active genes within a community, and measures the changes in the level of expression due to modified environmental conditions (Mukherjee and Reddy, 2020). Extracted RNAs are reverse-transcribed into cDNA and sequenced, as in metagenomics studies. Pioneering studies aiming to identify expressed genes in environmental samples date back to 2005 and represent the dawn of metatranscriptomics (Prosser, 2015). As in metagenomics studies, reads are either mapped to a reference genome or *de novo* assembled into contigs and scaffolds (Prosser, 2015). Several advantages of metatranscriptomics can be highlighted. This approach characterizes unknown microorganisms, as metagenomics does. It is complementary to metagenomics in that it provides a snapshot of expressed genes (as only coding sequences are obtained). Thus, the active microbiome is explored to understand how patterns of gene expression change following biotic and

abiotic stresses and various environmental disturbances. Yet, although it is easy to recover large quantities of total RNA from a pure bacterial culture (tens of micrograms *per* extraction), this is rather difficult in soil samples (Wang et al., 2012). According to reports, the extraction yields of soil RNA range from tens of nanograms to several micrograms *per* gram of soil. Such a range could be explained by the amount of living microorganisms in the soil samples, by contamination by humic substances or by RNA loss during purification (Wang et al., 2012). The technical issues associated with metatranscriptomics are similar to those associated with metagenomics, but with further potential biases introduced during rRNA depletion (as rRNAs can represent more than 80% of total RNAs), the construction of cDNA libraries, and the required rapid inactivation of samples to prevent mRNA turnover (Prosser, 2015).

As metatranscriptomics approaches can be complicated to apply across the whole food value chain, only specific compartments have been studied. For example, the microbiota of various Swiss-type cheeses were analyzed during ripening (Duru et al., 2018). Samples were collected from three cheeses at two time points – during warm room ripening and cold room ripening. The resulting data supported a better understanding of the flavor-forming mechanism, i.e., the succession of up-regulated and down-regulated pathways involving microbial species during warm and cold ripening of the cheese. Such studies are a first step toward improving the use of food microbiomes in terms of flavor, quality and security (Afshari et al., 2020).

Characterization of Microbial Community Functionality and Activity: Meta-Proteomics and Meta-Metabolomics

Meta-proteomics and meta-metabolomics are devoted to the description of microbial functionality and activity. Therefore, they are the functional complement of studies focused on DNA and/or RNA. Extracellular enzymes synthesized by microbial communities contribute to various ecosystem services such as organic matter decomposition or nutrient cycling (Van Emon, 2016). For this purpose, meta-proteomics characterizes the entire protein content of an environmental microbiome at a given location and time. Meta-metabolomics is dedicated to the comprehensive analysis of all metabolites (i.e., small molecules released by microorganisms into the environment) contained in a sample. Due to its large potential for providing a link between functional and phylogenetic information on soil microbial communities, there has been growing interest in the application of meta-proteomics in soil ecology to study microbe-driven ecosystem functions (Keiblinger et al., 2016). Bacteria, fungi and protists indeed excrete a wide range of volatile organic compounds to interact and communicate with each other, promote crop growth or modulate plant defense (Schulz-Bohm et al., 2017; Poveda, 2021).

Both approaches use common methodologies to extract proteins or metabolites from soil matrices. Direct extraction

can be applied, using physical, chemical or mechanical lysis of cells, leading to comprehensive protein (or metabolite) recovery from soil microorganisms. For the choice of the cell lysis method, the target proteins and soil texture should be considered (Keiblinger et al., 2016). This direct approach is often complex due to the presence of other organic compounds co-extracted with proteins or metabolites. These co-extracted products can also complexify the analysis steps. Moreover, the choice of the extraction method can clearly impact the results of the meta-metabolomics approach. Another solution consists in separating microbial cells from the soil matrix prior to extraction, e.g., by density gradient centrifugation (Keiblinger et al., 2016). This indirect extraction reduces the problems caused by interfering and coextracted substances. However, all proteins, enzymes and metabolites excreted by microorganisms are ignored.

Generating meta-proteomics and meta-metabolomics data differs significantly from generating metagenomics and metatranscriptomics data, which rely heavily on sequencing (Speda et al., 2017). Concerning meta-proteomics, the complexity of environmental samples still outstrips the capabilities of mass spectrometry approaches (Keiblinger et al., 2016). Thus, protein separation is mandatory to reduce sample complexity before mass spectrometry analysis. 2D gel-based protein separation methods prior to enzymatic digestion can be successfully employed, but these methods have major drawbacks, particularly regarding the analysis of proteins with extreme molecular weights, isoelectric points or hydrophobicity values. Gel-free approaches include different protein extraction procedures, followed by digestion into peptides. Peptides are further separated by reversed-phase liquid chromatography or 2D chromatographic separation using strong cation exchange chromatography in combination with reversed-phase liquid chromatography. Then, peptides are analyzed by mass spectrometry, the measured spectra are compared with theoretical spectra of peptides from a database. Due to its high efficiency and automation, this method is the main strategy for protein identification, quantification and detection. However, only protein sequences represented in the database can be identified.

Identifying and quantifying metabolites is typically carried out using a combination of chromatography techniques (i.e., liquid chromatography, gas chromatography) and detection methods, such as mass spectrometry, and nuclear magnetic resonance (Aguilar-Pulido et al., 2016b). These technologies produce spectra consisting of patterns of peaks that allow metabolite identification and quantification, using comparisons with spectral databases like BioMagResBank, the MassBank, or the Golm Metabolome Database (Aguilar-Pulido et al., 2016b). Prior to this analysis, denoising and peak-picking processes are essential to improve the quality of the treated data.

Data storage and management are a major issue for both approaches because the generated data take valuable space and are barely standardized (Keiblinger et al., 2016; Aguilar-Pulido et al., 2016b). Moreover, meta-proteomics and meta-metabolomics are relatively costly in terms of equipment, and need a high degree of scientific knowledge to compute, analyze and treat

the data. But, contrary to other meta-omics approaches, meta-proteomics and meta-metabolomics are the most direct methods for identifying the functionalities of a natural environment because they work as semi-quantitative methods. Moreover, as these approaches are very sensitive, they can be considered as a good bioindicator of soil quality. Due to their large potential for providing precise information on soil microbial communities, meta-proteomics and meta-metabolomics are currently most advanced among meta-omics approaches.

In parallel to metatranscriptomics, meta-proteomics and meta-metabolomics are generally applied only for specific compartments of food value chains (soils, plants, or transformed products). For example, the composition of soybean leaf metabolites was investigated using metabolomics (Yun et al., 2018) and linked with the soil physicochemical properties. The authors characterized leaf metabolites of 40 samples harvested at two growth stages and from two contrasting sites, with ten replicates *per* condition. They concluded that the soybean leaf metabolome is clearly dependent on the geographical area and influenced by environmental factors such as the soil properties. However, recent studies are a starting point for further, more detailed studies of more compartments of food chains (Mattarozzi et al., 2020). These authors applied meta-proteomics on rhizosphere soils to have a better understanding of the impact of biostimulants on maize seeds. They demonstrated that biostimulants increased the activity of the bacterial community to different extents, and created a permissive and nurturing rhizosphere by stimulating microbial metabolism. As a result, plant growth and root elongation were promoted, and the activity of the bacterial community – especially of species beneficial to plant growth – was increased, without the composition of the microbiota being modified.

CONTRIBUTION OF META-OMICS TO SOIL QUALITY INDICATORS

Issues and Obstacles to Making Meta-Omics Usable for Operational Bioindicators

As previously detailed, meta-omics can be used to characterize the soil microbial communities and evaluate the soil biological quality, but environmental heterogeneity can complexify the analyses. For microbial communities, environmental heterogeneity provides a multitude of habitats at the microscale and macroscale (Fierer, 2017; Karimi et al., 2020). Moreover, soil physico-chemical differences across fields, landscapes, regions and/or continents create an extraordinary pedodiversity that complexifies the study and the survey of soil microbial communities (Hermans et al., 2019). To ensure robust study strategies and avoid the effects of storage time and conditions on the microbial community composition, several international standardized methods (like ISO standards) have been developed to recommend the most accurate sampling method (composite,


stratified, systematic, random) and storage conditions of soil samples.

Along with the application of meta-omics methods to characterize microbial diversity in soil samples, several criteria must be verified to lead to an operational bioindicator (Figure 6). First, the repeatability, sensitivity and reproducibility levels of the method must be informed and controlled. Moreover, standard operating procedures must be created, evaluated, and tested by independent laboratories (Terrat et al., 2015). The chosen meta-omics procedure must be easily usable as routine analysis (Supplementary Table 1).

The understanding and interpretation of any bioindicator of soil quality needs a repository to diagnose the biological status of soils (Lemanceau et al., 2014; Hermans et al., 2017; Jansson and Hofmockel, 2018). Such repositories are used to evaluate whether the measured values are within the range of variations considered as “normal operating range,” namely for a given soil type and land use. Data rely on prior descriptive biodiversity studies, the determination of the relationships between microbial abundance and diversity, taxonomic abundance, microbial activities and functioning, and environmental filters (soil, physicochemical, meteorological and spatial parameters). This knowledge of microbial biogeography needs studies at a large spatial scale. A significant number of soil microbial surveys are listed at the national or territory scale (Figure 7). Some of these studies aim to monitor soil quality thanks to repositories, and propose modern and operational microbial indicators (Figure 7; Rutgers et al., 2009; Griffiths et al., 2011; Powell et al., 2015; Bissett et al., 2016; Horrigue et al., 2016; Terrat et al., 2017; Tresch et al., 2018; George et al., 2019; Ji et al., 2019; Norris et al., 2020). Some are used to measure the impact of land use changes combined with pedoclimatic conditions (Horrigue et al., 2016; Terrat et al., 2017), or of farming practices (Norris et al., 2020) or to evaluate the quality of urban soils (Tresch et al., 2018). The main measured indicators are microbial biomass, microbial diversity (16S, 18S rDNA and ITS markers) using metabarcoding, and more rarely the microbial functional potential through metagenomics (Bissett et al., 2016; Norris et al., 2020). Other research works have sought to understand the spatial distribution and the main environmental drivers, but can be considered as microbial monitoring frameworks for practical conservation of soils (Toju et al., 2016; Hermans et al., 2017; Coller et al., 2019; Jiao and Lu, 2020; Tedersoo et al., 2020; Figure 7).

Examples Using Meta-Omics to Assess Soil Quality as a Part of the Food Value Chain

The extensive use of meta-omics approaches to better characterize soil microbial communities has clearly improved our knowledge on soil functioning. However, the integration of -omics data to understand and evaluate all the compartments of the food value chain remain scarce and only some recent studies used it to deal with the microbiological risk assessment or the




agroecological transition, with the objective of a high food chain quality.

Impact of Agricultural Practices on Food Security

Soil fertilization with animal-based organic products, such as manure, can cause human pathogens to develop in the soil (Berge et al., 2009; Semenov et al., 2009). One of the greatest challenges for microbiologists is the exploitation of -omics data in microbiological risk assessment (Cocolin et al., 2018). To date, few studies using metagenomics have assessed the pathogenic status of amended soils (Fang et al., 2015;

Meneghine et al., 2017). The most dominant human pathogens detected in soil after a long-term application of chicken manure would be *Mycobacterium tuberculosis* and *Mycobacterium ulcerans* (Fang et al., 2015). The study of (Meneghine et al., 2017) used whole metagenomics sequencing approaches to investigate the taxonomic and functional profiles of microbial communities in soil amended with organic compost made from animal carcasses. Bacterial pathogens generally affiliated with fecal contamination were found in low abundance in the soil (less than 0.4% of the total identified bacterial genera), and no toxigeny-related gene or fecal contaminant was detected. More recently, virulence genes and ARGs were searched



for in long-term manure-amended greenhouse soils using metagenomics (Zhang et al., 2020). Manure application led to the accumulation of such virulence and ARGs in soils, and accumulation increased with the frequency of manure application. Such results highlight a risk of widespread ARGs and virulence genes in amended agricultural soils, that can be regulated by mitigating manure application frequency or by pre-treating manure.

From Soil Microbiology to Food Quality at the Territorial Scale

The agroecological transition praised at a global scale, but it operates at local scales where particular environmental,

social and economic constraints weigh on agricultural production (Schmitt et al., 2017). Based on this observation, projects are now developing that target focused areas and take into account the whole food value chain, from soils to food quality and consumption, including economic aspects (Magrini et al., 2019).

In France, products with labels of quality of origin (e.g., Protected Designation of Origin, aka. PDO; Protected Geographic Indication; “Label Rouge”) are good examples of food value chains in which the quality or the specificity of the products are related to the specificities of the “terroirs” of production, including the local soil properties. Production specifications have long been established independently of environmental issues, hence potential soil degradation. The

question of the relationship between the soil ecological quality and the quality of agricultural products has been regularly suggested in recent years, but it still needs objective scientific demonstrations. The IFEP project was set up to bring new scientific knowledge in this field by focusing on the relationship between soil and product specificity for Comté PDO cheese. It aims at investigating the microbial communities along the continuum between the soil, the grass phyllosphere, cow teats and milk, using a metabarcoding approach and across a network of 44 dairy farms (Chemidlin Prévost-Bouré et al., 2021). The results highlighted strong and significant links between the four compartments of this continuum. These links were modulated by environmental factors: soil pH, plant diversity and elevation, but also farming practices, in particular organic fertilization, cattle density and cow-teat care. Thanks to metabarcoding, the IFEP project provided a proof-of-concept of the importance of the soil microbial quality in the definition of “terroirs,” and more largely in the food value chain and its sustainability through microbial transfers from agroecosystems to final products. The next step will be to investigate to what extent integrating soil microbiological quality may improve the sustainability of the food production models, with projects extending this approach to other products and going further up the food value chain. To answer this question, an approach similar to the Living-lab appeared to be most promising and relevant, due to its numerous advantages for innovation in agriculture (Lacoste et al., 2022).

The “Dijon, sustainable food supply 2030” project is another example of the integration of soil quality in the food value chain at a territorial scale. This project is an ambitious and innovative ten-year plan aimed at reaching a fully sustainable agri-food model in the Dijon Métropole territory (Burgundy, France). Even though not obvious, metabarcoding of the soil DNA represents an important asset for this project. It will be implemented on a sampling of 600 points covering the entire 3,500 km² of Dijon Métropole’s urban and agricultural territories, and will lead to several outputs. From a scientific point of view, the implementation of metabarcoding on a territorial scale will provide more insights into the drivers of the spatial distribution and regulation of the soil microbial diversity. From a more operational point of view, it will lead to building up the first territorial repository of the soil microbial quality in terms of microbial biomass and biodiversity. Such a repository will serve to implement a robust and sensitive diagnosis of the soil microbial quality by the actors and stakeholders involved in the use of soils (stakeholders of biomonitoring programs, farmers, land managers, agri-food industries, and consumers). This is a pre-requisite for combining food and the environmental quality of future farming systems developed in the context of the agroecological transition. Indicators and associated repositories will also be used to implement innovative labels for agricultural products, based on performance obligations rather than just best endeavors obligations. Such labels will inform the consumers about the effective environmental fingerprint of the agricultural

products. They will also help move toward the development of local food systems. In addition, based on diagnosis at the territorial scale, the most sustainable farming systems and practices will be identified and communicated to the stakeholders involved in the use of soils, whether urban or rural. Soil quality indicators obtained from metabarcoding will also provide bases for soil value estimates integrating the intrinsic ecological state, which has not been taken into account to date. Ultimately, the implications of this shift of paradigm will be huge in terms of soil conservation, legal terms and economic aspects. Altogether, this project illustrates to what extent metabarcoding can be a powerful tool to help in the transition of urban and rural spaces toward more sustainable systems. As a pilot project, it is aimed at being replicated in other territories in order to supply collectivities and soil users with tools for measuring the impact of their activities on the soil quality and promoting sustainable systems.

ONGOING AND FUTURE DEVELOPMENTS


Recent Advanced Technologies

New technological developments frequently emerge in microbiology. Adapted from the Lawrence Livermore Microbial Detection Array (LLMDA), the Axiom Microbiome Array (AMA) is a high-throughput platform that identifies species from environmental samples. This microarray contains 1.38 million DNA probes that can detect just over 12,000 species of viruses, archaea and bacteria, fungi, and protozoa, and probably represents the most thorough method (Thissen et al., 2019).

New platforms devoted to third-generation sequencing have emerged. They provide long reads and avoid PCR steps for some technologies (Tedesoo et al., 2021). Although these technologies are increasingly used in microbial ecology research, they are not yet available for the microbial diagnosis of soils. Nevertheless, they have proved useful to characterize microbial communities in a context of diagnosis in plant pathology (Tedesoo et al., 2019). The small and large subunits of the rRNA operon can be sequenced in full length (Tedesoo et al., 2018), improving the species-level identification of prokaryote and eukaryote organisms (Benítez-Páez et al., 2016; Singer et al., 2016) and *in fine* diagnosis. The feasibility of the method was demonstrated by the Oxford Nanopore MinION™ system and PacBio sequencing with environmental samples for metabarcoding (Schloss et al., 2016; Kerkhof, 2021) and metagenomics (Van Goethem et al., 2021) studies. Real-time DNA sequencing coupled with bioinformatic analysis also appears very promising to carry out a rapid assessment of biodiversity directly in the field and might represent a reliable bio-monitoring tool in the years to come (Pomerantz et al., 2018; Krehenwinkel et al., 2019; Maestri et al., 2019; Albrecht et al., 2020).

Recent meta-omics approaches are essential to better understand the soil microbiome, but it would be interesting

to also explore the microorganisms of relevance for improving the soil quality and food safety. Although significant improvements have been made in culturing so-far unculturable microorganisms, finding an appropriate culture medium remains difficult. High-throughput culturing platforms have emerged to isolate and screen a large number of microbial cells (Mafla-Endara et al., 2021; Trivedi et al., 2021). The ability to isolate single cells using microfluidics coupled with technologies amplifying genomic DNA from single cells will revolutionize the study of unculturable species and of microheterogeneity within species (Chen et al., 2021). As recently stated, microfluidics has a great potential for experimenting on the soil microbiome so as to determine the mechanisms underpinning specific microbial metabolic interactions and to understand and predict the influence of environmental gradients on specific microbial functions with precise spatiotemporal control (Jansson and Hofmockel, 2018). The latest data indicated that droplet-based microfluidic systems compartmentalize single microorganisms in droplets, detect and sort the droplets at a rate of up to 1,000 per second and screen ~1 million microorganisms per hour (Watterson et al., 2020).



Old Data for Future Prospects, New Perspectives

Over the last decade, multiple tools or databases have been developed to maximize the use of meta-omics results. Major advances have been made in metabarcoding research, and have provided functional information (Djemiel et al., 2022). Functional inference and functional trait assignments can be carried out in addition to the characterization of microbial diversity. Functional inference based on phylogenetic models and genomic data predicts putative bacterial and fungal functions (Douglas et al., 2020). One of the advantages is to go beyond the metagenomics approach while saving time and money. However, the major limitation is that the robustness of the results is strongly linked to amplicon resolution (corresponding to metabarcoding) and the availability of microbial reference genomes (corresponding to inference). Further tools and databases can be used to supplement microbial diversity results with trait and character data (Nguyen et al., 2016; Pölme et al., 2020). For example, based on taxonomic classification, fungal genera/species can be associated with their lifestyles, body types or habitats. Another alternative way to evaluate the quality and functioning of soil microbial ecosystems is biotic interaction network analysis (Barberán et al., 2012; Layeghifard et al., 2017). Based on the analysis of taxonomic co-abundance patterns, the topology of each network highlights the relationship between taxa – positive in the case of co-occurrence, or negative in the case of co-exclusion (Karimi et al., 2017). Exploring the keystone taxa or guilds highly connected whatever their abundance will provide a complementary approach and detect taxa with a crucial role in the soil microbiota structure and functioning (Banerjee et al., 2018), which can be modified by soil disturbances linked to agricultural intensification (Banerjee et al., 2019; Karimi et al., 2019).

CONCLUSION

Global awareness is required for a greater consideration of microbial indicators for monitoring soil biodiversity and achieving the ambition of a sustainable food value chain. This involves a wide-scale collaboration of governments, local authorities, stakeholders of bio-monitoring programs, farmers, land managers, agri-food industries, and consumers (Lehmann et al., 2020; Fierer et al., 2021; **Figure 8**). Microbial indicators for soil quality monitoring by scientific researchers (Fierer et al., 2021) and routine use by field professionals (**Figure 8**) need to be developed and validated. Although studying microbial diversity used to be difficult owing to technical constraints, a broad range of approaches have emerged in the past few years or decades (Yang et al., 2020). Meta-omics techniques are already being used to study microbial communities along the food production chain, but insufficiently integrate the soil microbiome (Cocolin et al., 2018; Yap et al., 2022). This holistic vision is essential to offer high-value bioindicators. Moreover,

robust repositories are badly needed for presently existing soil monitoring networks (Yang et al., 2020). In light of these elements, meta-omics has provided and will keep on providing modern microbial indicators for monitoring the soil biodiversity and securing food production and quality, and the related environmental footprint with the creation of labels for the consumers. This ambition to achieve sustainable food systems also applies to urban environments (Vargas-Hernández, 2020).

AUTHOR CONTRIBUTIONS

CD and ST conceptualized the manuscript and drafted the manuscript with contributions from SD, BK, AC, WH, ArB, AIB, SS-B, P-AM, NCP-B, and LR. All authors contributed to the article and approved the submitted version.

REFERENCES

- Abot, A., Arnal, G., Auer, L., Lazuka, A., Labourdette, D., Lamarre, S., et al. (2016). CAZyChip: dynamic assessment of exploration of glycoside hydrolases in microbial ecosystems. *BMC Genomics* 17:671. doi: 10.1186/s12864-016-2988-4
- Afshari, R., Pillidge, C. J., Dias, D. A., Osborn, A. M., and Gill, H. (2020). Cheesomics: the future pathway to understanding cheese flavour and quality. *Crit. Rev. Food Sci. Nutr.* 60, 33–47. doi: 10.1080/10408398.2018.1512471
- Aguiar-Pulido, V., Huang, W., Suarez-ulloa, V., Cickovski, T., Mathee, K., and Narasimhan, G. (2016a). Approaches for microbiome analysis. *Evol. Bioinforma.* 12, 5–16. doi: 10.4137/EBO.S36436.TYPE
- Aguiar-Pulido, V., Huang, W., Suarez-Ulloa, V., Cickovski, T., Mathee, K., and Narasimhan, G. (2016b). Metagenomics, metatranscriptomics, and metabolomics approaches for microbiome analysis. *Evol. Bioinforma.* 12, 5–16. doi: 10.4137/EBO.S36436
- Albrecht, B., Bağcı, C., and Huson, D. H. (2020). MAIRA- real-time taxonomic and functional analysis of long reads on a laptop. *BMC Bioinformatics* 21:390. doi: 10.1186/s12859-020-03684-2
- Astudillo-García, C., Hermans, S. M., Stevenson, B., Buckley, H. L., and Lear, G. (2019). Microbial assemblages and bioindicators as proxies for ecosystem health status: potential and limitations. *Appl. Microbiol. Biotechnol.* 103, 6407–6421. doi: 10.1007/s00253-019-09963-0
- Banerjee, S., Schlaeppli, K., and van der Heijden, M. G. A. (2018). Keystone taxa as drivers of microbiome structure and functioning. *Nat. Rev. Microbiol.* 16, 567–576. doi: 10.1038/s41579-018-0024-1
- Banerjee, S., Walder, F., Büchi, L., Meyer, M., Held, A. Y., Gattinger, A., et al. (2019). Agricultural intensification reduces microbial network complexity and the abundance of keystone taxa in roots. *ISME J.* 13, 1722–1736. doi: 10.1038/s41396-019-0383-2
- Banos, S., Lentendu, G., Kopf, A., Wubet, T., Glöckner, F. O., and Reich, M. (2018). A comprehensive fungi-specific 18S rRNA gene sequence primer toolkit suited for diverse research issues and sequencing platforms. *BMC Microbiol.* 18:190. doi: 10.1186/s12866-018-1331-4
- Barberán, A., Bates, S. T., Casamayor, E. O., and Fierer, N. (2012). Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J.* 6, 343–351. doi: 10.1038/ismej.2011.119
- Barbi, F., Bragalini, C., Vallon, L., Prudent, E., Dubost, A., Fraissinet-Tachet, L., et al. (2014). PCR primers to study the diversity of expressed fungal genes encoding lignocellulolytic enzymes in soils using high-throughput sequencing. *PLoS One* 9:e116264. doi: 10.1371/journal.pone.0116264
- Bardgett, R. D., and Van Der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature* 515, 505–511. doi: 10.1038/nature13855
- Béné, C., Prager, S. D., Achicanoy, H. A. E., Toro, P. A., Lamotte, L., Bonilla, C., et al. (2019). Global map and indicators of food system sustainability. *Sci. Data* 6:279. doi: 10.1038/s41597-019-0301-5

ACKNOWLEDGMENTS

The authors thank Alexandrine Thore for graphical design in Figures 1, 3, 6, and 8. They also thank Annie Buchwalter for correcting and improving English language in the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2022.889788/full#supplementary-material>

Supplementary Table 1 | Overview of the various meta-omics approaches and their operationality for soil quality biomonitoring. R: In research phase, P: Provided by agri-food industries, XXX: Adapted technique with a very good data / price ratio, XX: Adapted technique with a moderate data / price ratio, X: Adapted technique with a low data/price ratio.

- Benítez-Páez, A., Portune, K. J., and Sanz, Y. (2016). Species-level resolution of 16S rRNA gene amplicons sequenced through the MinION™ portable nanopore sequencer. *Gigascience* 5, 1–9. doi: 10.1186/s13742-016-0111-z
- Berge, A. C. B., Glanville, T. D., Millner, P. D., and Klingborg, D. J. (2009). Methods and microbial risks associated with composting of animal carcasses in the United States. *J. Am. Vet. Med. Assoc.* 234, 47–56. doi: 10.2460/javma.234.1.47
- Bissett, A., Fitzgerald, A., Meintjes, T., Mele, P. M., Reith, F., Dennis, P. G., et al. (2016). Introducing BASE: The biomes of Australian soil environments soil microbial diversity database. *Gigascience* 5:s13742-016. doi: 10.1186/s13742-016-0126-5
- Bloem, J., Hopkins, D. W., and Benedetti, A. (2005). *Microbiological Methods for Assessing Soil Quality*. United Kingdom: CABI.
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* 37, 852–857. doi: 10.1038/s41587-019-0209-9
- Bukin, Y. S., Galachyants, Y. P., Morozov, I. V., Bukin, S. V., Zakharenko, A. S., and Zemskaya, T. I. (2019). The effect of 16S rRNA region choice on bacterial community metabarcoding results. *Sci. Data* 6, 1–14. doi: 10.1038/sdata.2019.7
- Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., De Goede, R., et al. (2018). Soil quality – A critical review. *Soil Biol. Biochem.* 120, 105–125. doi: 10.1016/j.soilbio.2018.01.030
- Castaño, C., Berlin, A., Brandström Durling, M., Ihrmark, K., Lindahl, B. D., Stenlid, J., et al. (2020). Optimized metabarcoding with Pacific biosciences enables semi-quantitative analysis of fungal communities. *New Phytol.* 228:nph.16731. doi: 10.1111/nph.16731
- Chemidlin Prévost-Bouré, N., Karimi, B., Sadet-Bourgeteau, S., Djemiel, C., Brie, M., Dumont, J., et al. (2021). Microbial transfers from permanent grassland ecosystems to milk in dairy farms in the Comté cheese area. *Sci. Rep.* 11:18144. doi: 10.1038/s41598-021-97373-6
- Chen, Q. L., Hu, H. W., He, Z. Y., Cui, L., Zhu, Y. G., and He, J. Z. (2021). Potential of indigenous crop microbiomes for sustainable agriculture. *Nat. Food* 2, 233–240. doi: 10.1038/s43016-021-00253-5
- Chesworth, W. (2007). *Encyclopedia of Soil Science*. Germany: Springer Science.
- Cocolin, L., Mataragas, M., Bourdichon, F., Doulgeraki, A., Pilet, M. F., Jagadeesan, B., et al. (2018). Next generation microbiological risk assessment meta-omics: The next need for integration. *Int. J. Food Microbiol.* 287, 10–17. doi: 10.1016/j.ijfoodmicro.2017.11.008
- Coller, E., Cestaro, A., Zanzotti, R., Bertoldi, D., Pindo, M., Larger, S., et al. (2019). Microbiome of vineyard soils is shaped by geography and management. *Microbiome* 7:140. doi: 10.1186/s40168-019-0758-7
- Cui, H., Wang, C., Gu, Z., Zhu, H., Fu, S., and Yao, Q. (2014). Evaluation of soil storage methods for soil microbial community using genetic and

- metabolic fingerprintings. *Eur. J. Soil Biol.* 63, 55–63. doi: 10.1016/j.ejsobi.2014.05.006
- Delmont, T. O., Robe, P., Clark, I., Simonet, P., and Vogel, T. M. (2011). Metagenomic comparison of direct and indirect soil DNA extraction approaches. *J. Microbiol. Methods* 86, 397–400. doi: 10.1016/j.mimet.2011.06.013
- Dickie, I. A., Boyer, S., Buckley, H., Duncan, R. P., Gardner, P., Hogg, I. D., et al. (2018). Towards robust and repeatable sampling methods in eDNA based studies. *Mol. Ecol. Resour.* 5, 940–952. doi: 10.1111/1755-0998.12907
- Djemiel, C., Dequiedt, S., Karimi, B., Cottin, A., Girier, T., El Djoudi, Y., et al. (2020a). BIOCOP-PIPE: a new user-friendly metabarcoding pipeline for the characterization of microbial diversity from 16S, 18S and 23S rRNA gene amplicons. *BMC Bioinformatics* 21:492. doi: 10.1186/s12859-020-03829-3
- Djemiel, C., Maron, P., Terrat, S., Dequiedt, S., Cottin, A., and Ranjard, L. (2022). Inferring microbiota functions from taxonomic genes: a review. *Gigascience* 11, 1–30. doi: 10.1093/gigascience/giab090
- Djemiel, C., Plassard, D., Terrat, S., Crouzet, O., Sauze, J., Mondy, S., et al. (2020b). μ green-db: a reference database for the 23S rRNA gene of eukaryotic plastids and cyanobacteria. *Sci. Rep.* 10, 1–11. doi: 10.1038/s41598-020-62555-1
- Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M., et al. (2020). PICRUSt2 for prediction of metagenome functions. *Nat. Biotechnol.* 38, 685–688. doi: 10.1038/s41587-020-0548-6
- Duru, I. C., Laine, P., Andreevskaya, M., Paulin, L., Kananen, S., Tynkynen, S., et al. (2018). Metagenomic and metatranscriptomic analysis of the microbial community in Swiss-type Maasdam cheese during ripening. *Int. J. Food Microbiol.* 281, 10–22. doi: 10.1016/j.ijfoodmicro.2018.05.017
- Escudé, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., et al. (2018). FROGS: find, rapidly, OTUs with galaxy solution. *Bioinformatics* 34, 1287–1294. doi: 10.1093/bioinformatics/btx791
- Evans, D. L., Quinton, J. N., Tye, A. M., Rodés, Á., Davies, J. A. C., Mudd, S. M., et al. (2019). Arable soil formation and erosion: a hillslope-based cosmogenic nuclide study in the United Kingdom. *Soil* 5, 253–263. doi: 10.5194/soil-5-253-2019
- Fang, H., Wang, H., Cai, L., and Yu, Y. (2015). Prevalence of antibiotic resistance genes and bacterial pathogens in long-term manured greenhouse soils As revealed by metagenomic survey. *Environ. Sci. Technol.* 49, 1095–1104. doi: 10.1021/es504157v
- FAO, ITPS, GSBi, and SCBD (2020). *State of Knowledge of Soil Biodiversity - Status, Challenges and Potentialities*. Rome, Italy: FAO. 618.
- FAOSTAT (2021). Food and Agriculture Organization of the United Nations (FAO). Available at: <https://www.fao.org/faostat/en/#home>
- Fierer, N. (2017). Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev. Microbiol.* 15, 579–590. doi: 10.1038/nrmicro.2017.87
- Fierer, N., Wood, S. A., and Bueno de Mesquita, C. P. (2021). How microbes can, and cannot, be used to assess soil health. *Soil Biol. Biochem.* 153:108111. doi: 10.1016/j.soilbio.2020.108111
- Frac, M., Hannula, S. E., Belka, M., and Jędrzycka, M. (2018). Fungal biodiversity and their role in soil health. *Front. Microbiol.* 9:707. doi: 10.3389/fmicb.2018.00707
- Friedrichsen, C. N., Daroub, S. H., Monroé, M. C., Stepp, J. R., and Gerber, S. (2019). Stakeholders' mental models of soil food value chain in the Everglades. *Geoderma* 343, 166–175. doi: 10.1016/j.geoderma.2019.02.037
- Geisen, S., Mitchell, E. A. D., Adl, S., Bonkowski, M., Dunthorn, M., Ekelund, E., et al. (2018). Soil protists: a fertile frontier in soil biology research. *FEMS Microbiol. Rev.* 42, 293–323. doi: 10.1093/femsre/fuy006
- Geisen, S., Wall, D. H., and van der Putten, W. H. (2019). Challenges and opportunities for soil biodiversity in the Anthropocene. *Curr. Biol.* 29, R1036–R1044. doi: 10.1016/j.cub.2019.08.007
- George, P. B. L., Lallias, D., Creer, S., Seaton, F. M., Kenny, J. G., Eccles, R. M., et al. (2019). Divergent national-scale trends of microbial and animal biodiversity revealed across diverse temperate soil ecosystems. *Nat. Commun.* 10, 1–11. doi: 10.1038/s41467-019-09031-1
- Goodwin, S., McPherson, J. D., and McCombie, W. R. (2016). Coming of age: ten years of next-generation sequencing technologies. *Nat. Rev. Genet.* 17, 333–351. doi: 10.1038/nrg.2016.49
- Griffiths, B. S., Faber, J., and Bloem, J. (2018). Applying soil health indicators to encourage sustainable soil use: The transition from scientific study to practical application. *Sustainability* 10:3021. doi: 10.3390/su10093021
- Griffiths, R. I., Thomson, B. C., James, P., Bell, T., Bailey, M., and Whiteley, A. S. (2011). The bacterial biogeography of British soils. *Environ. Microbiol.* 13, 1642–1654. doi: 10.1111/j.1462-2920.2011.02480.x
- Gutiérrez-Aguirre, I., Rački, N., Dreo, T., and Ravnika, M. (2015). “Droplet digital PCR for absolute quantification of pathogens,” in *Plant Pathology* (New York, NY: Humana Press), 331–347.
- Hadziavdic, K., Lekang, K., Lanzen, A., Jonassen, I., Thompson, E. M., and Troedsson, C. (2014). Characterization of the 18S rRNA gene for designing universal eukaryote specific primers. *PLoS One* 9:e87624. doi: 10.1371/journal.pone.0087624
- Hambraeus, G., Von Wachenfeldt, C., and Hederstedt, L. (2003). Genome-wide survey of mRNA half-lives in *Bacillus subtilis* identifies extremely stable mRNAs. *Mol. Gen. Genomics.* 269, 706–714. doi: 10.1007/s00438-003-0883-6
- Hartemink, A. E., Zhang, Y., Bockheim, J. G., Curi, N., Silva, S. H. G., Grauer-Gray, J., et al. (2020). Soil horizon variation: A review. *Adv. Agron.* 160, 125–185. doi: 10.1016/bs.agron.2019.10.003
- Hayatsu, M., Tago, K., and Saito, M. (2008). Various players in the nitrogen cycle: diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Sci. Plant Nutr.* 54, 33–45. doi: 10.1111/j.1747-0765.2007.00195.x
- He, Z., Gentry, T. J., Schadt, C. W., Wu, L., Liebich, J., Chong, S. C., et al. (2007). GeoChip: A comprehensive microarray for investigating biogeochemical, ecological and environmental processes. *ISME J.* 1, 67–77. doi: 10.1038/ismej.2007.2
- Hermans, S. M., Buckley, H. L., Case, B. S., Curran-Cournane, F., Taylor, M., and Lear, G. (2017). Bacteria as emerging indicators of soil condition. *Appl. Environ. Microbiol.* 83:e02826. doi: 10.1128/AEM.02826-16
- Hermans, S. M., Buckley, H. L., and Lear, G. (2019). Perspectives on the impact of sampling design and intensity on soil microbial diversity estimates. *Front. Microbiol.* 10:1820. doi: 10.3389/fmicb.2019.01820
- Hofstetter, V., Buyck, B., Eyssartier, G., Schnee, S., and Gindro, K. (2019). The unbearable lightness of sequenced-based identification. *Fungal Divers.* 96, 243–284. doi: 10.1007/s13225-019-00428-3
- Holmes, D. E., Nevin, K. P., and Lovley, D. R. (2004). Comparison of 16S rRNA, nifD, recA, gyrB, rpoB and fusA genes within the family Geobacteraceae fam. Nov. *Int. J. Syst. Evol. Microbiol.* 54, 1591–1599. doi: 10.1099/ijs.0.02958-0
- Horrigue, W., Dequiedt, S., Chemidlin Prévost-Bouré, N., Jolivet, C., Saby, N. P. A., Arrouays, D., et al. (2016). Predictive model of soil molecular microbial biomass. *Ecol. Indic.* 64, 203–211. doi: 10.1016/j.ecolind.2015.12.004
- Jansson, J. K., and Hofmockel, K. S. (2018). The soil microbiome — from metagenomics to metaphenomics. *Curr. Opin. Microbiol.* 43, 162–168. doi: 10.1016/j.mib.2018.01.013
- Ji, N., Gao, C., Sandel, B., Zheng, Y., Chen, L., Wu, B., et al. (2019). Late Quaternary climate change explains soil fungal community composition rather than fungal richness in forest ecosystems. *Ecol. Evol.* 9, 6678–6692. doi: 10.1002/ece3.5247
- Jiao, S., and Lu, Y. (2020). Abundant fungi adapt to broader environmental gradients than rare fungi in agricultural fields. *Glob. Chang. Biol.* 26, 4506–4520. doi: 10.1111/gcb.15130
- Karimi, B., Dequiedt, S., Terrat, S., Jolivet, C., Arrouays, D., Wincker, P., et al. (2019). Biogeography of soil bacterial networks along a gradient of cropping intensity. *Sci. Rep.* 9, 1–10. doi: 10.1038/s41598-019-40422-y
- Karimi, B., Maron, P. A., Bouré, C. P., Bernard, N., Gilbert, D., and Ranjard, L. (2017). Microbial diversity and ecological networks as indicators of environmental quality. *Environ. Chem. Lett.* 15, 265–281. doi: 10.1007/s10311-017-0614-6
- Karimi, B., Villerd, J., Dequiedt, S., Terrat, S., Chemidlin-Prévost Bouré, N., Djemiel, C., et al. (2020). Biogeography of soil microbial habitats across France. *Glob. Biogeogr.* 29, 1399–1411. doi: 10.1111/geb.13118
- Keiblinger, K. M., Fuchs, S., Zechmeister-Boltenstern, S., and Riedel, K. (2016). Soil and leaf litter metaproteomics-A brief guideline from sampling to understanding. *FEMS Microbiol. Ecol.* 92, 1–18. doi: 10.1093/femsec/iw180
- Kerkhof, L. J. (2021). Is Oxford Nanopore sequencing ready for analyzing complex microbiomes? *FEMS Microbiol. Ecol.* 97:fiab001. doi: 10.1093/femsec/fiab001
- Kieser, S., Brown, J., Zdobnov, E. M., Trajkovski, M., and McCue, L. A. (2020). ATLAS: a Snakemake workflow for assembly, annotation, and genomic binning of metagenome sequence data. *BMC Bioinform.* 21:257. doi: 10.1186/s12859-020-03585-4
- Kopittke, P. M., Berhe, A. A., Carrillo, Y., Cavagnaro, T. R., Chen, D., Chen, Q.-L., et al. (2022). Ensuring planetary survival: the centrality of organic carbon

- in balancing the multifunctional nature of soils. *Crit. Rev. Environ. Sci. Technol.* 1–17. doi: 10.1080/10643389.2021.2024484
- Kopittke, P. M., Menzies, N. W., Wang, P., McKenna, B. A., and Lombi, E. (2019). Soil and the intensification of agriculture for global food security. *Environ. Int.* 132:105078. doi: 10.1016/j.envint.2019.105078
- Krehenwinkel, H., Pomerantz, A., and Prost, S. (2019). Genetic biomonitoring and biodiversity assessment using portable sequencing technologies: current uses and future directions. *Genes* 10:858. doi: 10.3390/genes10110858
- Kultima, J. R., Coelho, L. P., Forslund, K., Huerta-Cepas, J., Li, S. S., Driessen, M., et al. (2016). MOCAT2: a metagenomic assembly, annotation and profiling framework. *Bioinformatics* 32, 2520–2523. doi: 10.1093/bioinformatics/btw183
- Lacoste, M., Cook, S., McNee, M., Gale, D., Ingram, J., Bellon-Maurel, V., et al. (2022). On-farm experimentation to transform global agriculture. *Nat. Food* 3, 11–18. doi: 10.1038/s43016-021-00424-4
- Layeghifard, M., Hwang, D. M., and Guttman, D. S. (2017). Disentangling interactions in the microbiome: A network perspective. *Trends Microbiol.* 25, 217–228. doi: 10.1016/j.tim.2016.11.008
- Lee, Y. J., Van Nostrand, J. D., Tu, Q., Lu, Z., Cheng, L., Yuan, T., et al. (2013). The PathoChip, a functional gene array for assessing pathogenic properties of diverse microbial communities. *ISME J.* 7, 1974–1984. doi: 10.1038/ismej.2013.88
- Lehmann, J., Bossio, D. A., Kögel-Knabner, I., and Rillig, M. C. (2020). The concept and future prospects of soil health. *Nat. Rev. Earth Environ.* 1, 544–553. doi: 10.1038/s43017-020-0080-8
- Lemanceau, P., Maron, P. A., Mazurier, S., Mougél, C., Pivato, B., Plassart, P., et al. (2014). Understanding and managing soil biodiversity: a major challenge in agroecology. *Agron. Sustain. Dev.* 35, 67–81. doi: 10.1007/s13593-014-0247-0
- Maestri, C., Paterno, F., and Garces, M. (2019). A rapid and accurate MinION-based workflow for tracking species biodiversity in the field. *Genes* 10:468. doi: 10.3390/genes10060468
- Mafla-Endara, P. M., Arellano-Caicedo, C., Aleklett, K., Pucetaite, M., Ohlsson, P., and Hammer, E. C. (2021). Microfluidic chips provide visual access to in situ soil ecology. *Commun. Biol.* 4:889. doi: 10.1038/s42003-021-02379-5
- Magrini, M.-B., Martin, G., Magne, M.-A., Duru, M., Couix, N., Hazard, L., et al. (2019). “Agroecological transition from farms to territorialised Agri-food systems: issues and drivers,” in *Agroecological Transitions: From Theory to Practice in Local Participatory Design*. eds. J. E. Bergez, E. Audouin and O. Therond (Cham: Springer International Publishing), 69–98.
- Maron, P.-A., Mougél, C., and Ranjard, L. (2011). Soil microbial diversity: methodological strategy, spatial overview and functional interest. *C. R. Biol.* 334, 403–411. doi: 10.1016/j.crv.2010.12.003
- Maron, P.-A., Sarr, A., Kaisermann, A., Lévêque, J., Mathieu, O., Guigüe, J., et al. (2018). High microbial diversity promotes soil ecosystem functioning. *Appl. Environ. Microbiol.* 84:e02738-17. doi: 10.1128/AEM.02738-17
- Martinez-Salgado, M. M., Gutierrez-Romero, V., Janssens, M., and Ortega-Blu, R. (2010). Biological soil quality indicators: a review. *Curr. Res. Technol. Educ. Top. Appl. Microbiol. Microb. Biotechnol.* 319–328.
- Martin-Laurent, F., Philippot, L., Hallet, S., Chaussod, R., Germon, J. C., Soulas, G., et al. (2001). DNA extraction from soils: old Bias for new microbial diversity analysis methods. *Appl. Environ. Microbiol.* 67, 2354–2359. doi: 10.1128/AEM.67.5.2354-2359.2001
- Mattarozzi, M., Di Zinno, J., Montanini, B., Manfredi, M., Marengo, E., Fornasier, F., et al. (2020). Biot stimulants applied to maize seeds modulate the enzymatic activity and metaproteome of the rhizosphere. *Appl. Soil Ecol.* 148:103480. doi: 10.1016/j.apsoil.2019.103480
- Mehle, N., and Dreo, T. (2019). “Quantitative analysis with droplet digital PCR,” in *Notes on the Greek Text of Genesis*. ed. J. W. Wevers (United States: SBL Press).
- Meneghini, A. K., Nielsen, S., Varani, A. M., Thomas, T., and Carareto Alves, L. M. (2017). Metagenomic analysis of soil and freshwater from zoo agricultural area with organic fertilization. *PLoS One* 12:e0190178. doi: 10.1371/journal.pone.0190178
- Meyer, F., Paarmann, D., D’Souza, M., Olson, R., Glass, E., Kubal, M., et al. (2008). The metagenomics RAST server – a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9:386. doi: 10.1186/1471-2105-9-386
- Mukherjee, A., and Reddy, M. S. (2020). Metatranscriptomics: an approach for retrieving novel eukaryotic genes from polluted and related environments. *Biotech* 10:71. doi: 10.1007/s13205-020-2057-1
- Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., et al. (2016). FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol.* 20, 241–248. doi: 10.1016/j.funeco.2015.06.006
- Nielsen, U. N., Wall, D. H., and Six, J. (2015). Soil biodiversity and the environment. *Annu. Rev. Environ. Resour.* 40, 63–90. doi: 10.1146/annurev-environ-102014-021257
- Nilsson, R. H., Anslan, S., Bahram, M., Wurzbacher, C., Baldrian, P., and Tedersoo, L. (2019a). Mycobiome diversity: high-throughput sequencing and identification of fungi. *Nat. Rev. Microbiol.* 17, 95–109. doi: 10.1038/s41579-018-0116-y
- Nilsson, R. H., Larsson, K.-H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., et al. (2019b). The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* 47, D259–D264. doi: 10.1093/nar/gky1022
- Nkongolo, K. K., and Narendrula-Kotha, R. (2020). Advances in monitoring soil microbial community dynamic and function. *J. Appl. Genet.* 61, 249–263. doi: 10.1007/s13353-020-00549-5
- Norris, C. E., Bean, G. M., Cappellazzi, S. B., Cope, M., Greub, K. L. H., Liptzin, D., et al. (2020). Introducing the north American project to evaluate soil health measurements. *Agron. J.* 112, 3195–3215. doi: 10.1002/agj2.20234
- Ogier, J.-C., Pages, S., Galan, M., Barret, M., and Gaudriault, S. (2019). rpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. *BMC Microbiol.* 19:171. doi: 10.1186/s12866-019-1546-z
- Oshiki, M., Segawa, T., and Ishii, S. (2018). Nitrogen cycle evaluation (NiCE) Chip for simultaneous analysis of multiple N cycle-associated genes. *Appl. Environ. Microbiol.* 84:e02615-17. doi: 10.1128/AEM.02615-17
- Pauvert, C., Buée, M., Laval, V., Edel-Hermann, V., Fauchery, L., Gautier, A., et al. (2019). Bioinformatics matters: The accuracy of plant and soil fungal community data is highly dependent on the metabarcoding pipeline. *Fungal Ecol.* 41, 23–33. doi: 10.1016/j.funeco.2019.03.005
- Phillips, H. R. P., Bach, E. M., Bartz, M. L. C., Bennett, J. M., Beugnon, R., Briones, M. J. I., et al. (2021). Global data on earthworm abundance, biomass, diversity and corresponding environmental properties. *Sci. Data* 8:136. doi: 10.1038/s41597-021-00912-z
- Poirier, S., Rué, O., Peguilhan, R., Coeuret, G., Zagorec, M., Champomier-Vergès, M. C., et al. (2018). Deciphering intra-species bacterial diversity of meat and seafood spoilage microbiota using gyrB amplicon sequencing: A comparative analysis with 16S rDNA V3-V4 amplicon sequencing. *PLoS One* 13:e0204629. doi: 10.1371/journal.pone.0204629
- Pôlme, S., Abarenkov, K., Henrik Nilsson, R., Lindahl, B. D., Clemmensen, K. E., Kausserud, H., et al. (2020). FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Divers.* 105, 1–16. doi: 10.1007/s13225-020-00466-2
- Pomerantz, A., Peñafiel, N., Arteaga, A., Bustamante, L., Pichardo, F., Coloma, L. A., et al. (2018). Real-time DNA barcoding in a rainforest using nanopore sequencing: opportunities for rapid biodiversity assessments and local capacity building. *Gigascience* 7, 1–14. doi: 10.1093/gigascience/giy033
- Porter, T. M., and Hajibabaei, M. (2018). Scaling up: A guide to high-throughput genomic approaches for biodiversity analysis. *Mol. Ecol.* 27, 313–338. doi: 10.1111/mec.14478
- Poveda, J. (2021). Beneficial effects of microbial volatile organic compounds (MVOs) in plants. *Appl. Soil Ecol.* 168:104118. doi: 10.1016/j.apsoil.2021.104118
- Powell, J. R., Karunaratne, S., Campbell, C. D., Yao, H., Robinson, L., and Singh, B. K. (2015). Deterministic processes vary during community assembly for ecologically dissimilar taxa. *Nat. Commun.* 6:8444. doi: 10.1038/ncomms9444
- Prosser, J. I. (2015). Dispersing misconceptions and identifying opportunities for the use of “omics” in soil microbial ecology. *Nat. Rev. Microbiol.* 13, 439–446. doi: 10.1038/nrmicro3468
- Prudent, M., Dequiedt, S., Sorin, C., Girodet, S., Nowak, V., Duc, G., et al. (2020). The diversity of soil microbial communities matters when legumes face drought. *Plant Cell Environ.* 43, 1023–1035. doi: 10.1111/pce.13712
- Pulleman, M., Creamer, R., Hamer, U., Helder, J., Pelosi, C., Pérès, G., et al. (2012). Soil biodiversity, biological indicators and soil ecosystem services—an overview of European approaches. *Curr. Opin. Environ. Sustain.* 4, 529–538. doi: 10.1016/j.cosust.2012.10.009
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA ribosomal RNA gene database project: improved data

- processing and web-based tools. *Nucleic Acids Res.* 41, D590–D596. doi: 10.1093/nar/gks1219
- Ranjard, L., Dequiedt, S., Lelievre, M., Maron, P. A., Mougel, C., Morin, F., et al. (2009). Platform GenoSol: a new tool for conserving and exploring soil microbial diversity. *Environ. Microbiol. Rep.* 1, 97–99. doi: 10.1111/j.1758-2229.2009.00023.x
- Rincon-Florez, V., Carvalhais, L., and Schenk, P. (2013). Culture-independent molecular tools for soil and rhizosphere microbiology. *Diversity* 5, 581–612. doi: 10.3390/d5030581
- Rutgers, M., Schouten, A. J., Bloem, J., Van Eekeren, N., De Goede, R. G. M., Jagersop Akkerhuis, G. A. J. M., et al. (2009). Biological measurements in a nationwide soil monitoring network. *Eur. J. Soil Sci.* 60, 820–832. doi: 10.1111/j.1365-2389.2009.01163.x
- Schloss, P. D., Jenior, M. L., Koupmpouras, C. C., Westcott, S. L., and Highlander, S. K. (2016). Sequencing 16S rRNA gene fragments using the PacBio SMRT DNA sequencing system. *PeerJ* 4:e1869. doi: 10.7717/peerj.1869
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541. doi: 10.1128/AEM.01541-09
- Schlöter, M., Nannipieri, P., Sørensen, S. J., and Van Elsland, J. D. (2018). Microbial indicators for soil quality. *Biol. Fertil. Soils* 54, 1–10. doi: 10.1007/s00374-017-1248-3
- Schmitt, E., Galli, F., Menozzi, D., Maye, D., Touzard, J.-M., Marescotti, A., et al. (2017). Comparing the sustainability of local and global food products in Europe. *J. Clean. Prod.* 165, 346–359. doi: 10.1016/j.jclepro.2017.07.039
- Schulz-Bohm, K., Geisen, S., Wubs, E. R. J., Song, C., de Boer, W., and Garbeva, P. (2017). The prey's scent – volatile organic compound mediated interactions between soil bacteria and their protist predators. *ISME J.* 11, 817–820. doi: 10.1038/ismej.2016.144
- Semenov, M. V. (2021). Metabarcoding and metagenomics in soil ecology research: achievements, challenges, and prospects. *Biol. Bull. Rev.* 11, 40–53. doi: 10.1134/S2079086421010084
- Semenov, A. V., Van Overbeek, L., and Van Bruggen, A. H. C. (2009). Percolation and survival of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar typhimurium in soil amended with contaminated dairy manure or slurry. *Appl. Environ. Microbiol.* 75, 3206–3215. doi: 10.1128/AEM.01791-08
- Sessitsch, A., Hackl, E., Wenzl, P., Kilian, A., Kostic, T., Stralis-Pavese, N., et al. (2006). Diagnostic microbial microarrays in soil ecology. *New Phytol.* 171, 719–736. doi: 10.1111/j.1469-8137.2006.01824.x
- Silver, W. L., Perez, T., Mayer, A., and Jones, A. R. (2021). The role of soil in the contribution of food and feed. *Philos. Trans. R. Soc. B Biol. Sci.* 376:20200181. doi: 10.1098/rstb.2020.0181
- Singer, E., Bushnell, B., Coleman-Derr, D., Bowman, B., Bowers, R. M., Levy, A., et al. (2016). High-resolution phylogenetic microbial community profiling. *ISME J.* 10, 1–13. doi: 10.1038/ismej.2015.249
- Speda, J., Jonsson, B.-H., Carlsson, U., and Karlsson, M. (2017). Metaproteomics-guided selection of targeted enzymes for bioprospecting of mixed microbial communities. *Biotechnol. Biofuels* 10:128. doi: 10.1186/s13068-017-0815-z
- Srivastava, N., Gupta, B., Gupta, S., Danquah, M. K., and Sarethy, I. P. (2019). “Analyzing functional microbial diversity,” in *Microbial diversity in the genomic era*. eds. S. Das and H.R. Dash (United States: Academic Press).
- Stefanini, I., and Cavalieri, D. (2018). Metagenomic approaches to investigate the contribution of the vineyard environment to the quality of wine fermentation: potentials and difficulties. *Front. Microbiol.* 9:991. doi: 10.3389/fmicb.2018.00991
- Swenson, T. L., Jenkins, S., Bowen, B. P., and Northen, T. R. (2015). Untargeted soil metabolomics methods for analysis of extractable organic matter. *Soil Biol. Biochem.* 80, 189–198. doi: 10.1016/j.soilbio.2014.10.007
- Tedersoo, L., Albertsen, M., Anslan, S., and Callahan, B. (2021). Perspectives and benefits of high-throughput long-read sequencing in microbial ecology. *Appl. Environ. Microbiol.* 87, 1–19. doi: 10.1128/AEM.00626-21
- Tedersoo, L., Anslan, S., Bahram, M., Drenkhan, R., Pritsch, K., Buegger, F., et al. (2020). Regional-scale in-depth analysis of soil fungal diversity reveals strong pH and plant species effects in northern Europe. *Front. Microbiol.* 11:1953. doi: 10.3389/fmicb.2020.01953
- Tedersoo, L., Drenkhan, R., Anslan, S., Morales-Rodriguez, C., and Cleary, M. (2019). High-throughput identification and diagnostics of pathogens and pests: overview and practical recommendations. *Mol. Ecol. Resour.* 19, 47–76. doi: 10.1111/1755-0998.12959
- Tedersoo, L., Tooming-Klunderud, A., and Anslan, S. (2018). PacBio metabarcoding of Fungi and other eukaryotes: errors, biases and perspectives. *New Phytol.* 217, 1370–1385. doi: 10.1111/nph.14776
- Terrat, S., Christen, R., Dequiedt, S., Lelievre, M., Nowak, V., Regnier, T., et al. (2012). Molecular biomass and MetaTaxogenomic assessment of soil microbial communities as influenced by soil DNA extraction procedure. *Microb. Biotechnol.* 5, 135–141. doi: 10.1111/j.1751-7915.2011.00307.x
- Terrat, S., Horryguez, W., Dequiedt, S., Saby, A., Lelièvre, M., Nowak, V., et al. (2017). Correction: mapping and predictive variations of soil bacterial richness across France. *PLoS One* 12:e0186766. doi: 10.1371/journal.pone.0186766
- Terrat, S., Plassart, P., Bourgeois, E., Ferreira, S., Dequiedt, S., Adele-Dit-De-Renseville, N., et al. (2015). Meta-barcoded evaluation of the ISO standard 11063 DNA extraction procedure to characterize soil bacterial and fungal community diversity and composition. *Microb. Biotechnol.* 8, 131–142. doi: 10.1111/1751-7915.12162
- Thiele-Bruhn, S., Schloter, M., Wilke, B. M., Beaudette, L. A., Martin-Laurent, F., Cheviron, N., et al. (2020). Identification of new microbial functional standards for soil quality assessment. *Soil* 6, 17–34. doi: 10.5194/soil-6-17-2020
- Thissen, J. B., Be, N. A., McLoughlin, K., Gardner, S., Rack, P. G., Shaper, M. H., et al. (2019). Axiom microbiome Array, the next generation microarray for high-throughput pathogen and microbiome analysis. *PLoS One* 14, 1–17. doi: 10.1371/journal.pone.0212045
- Toju, H., Kishida, O., Katayama, N., and Takagi, K. (2016). Networks depicting the fine-scale co-occurrences of fungi in soil horizons. *PLoS One* 11:e0165987. doi: 10.1371/journal.pone.0165987
- Tresch, S., Moretti, M., Le Bayon, R.-C., Mäder, P., Zanetta, A., Frey, D., et al. (2018). Urban soil quality assessment—A comprehensive Case study dataset of urban garden soils. *Front. Environ. Sci.* 6, 1–5. doi: 10.3389/fenvs.2018.00136
- Trivedi, P., Mattupalli, C., Eversole, K., and Leach, J. E. (2021). Enabling sustainable agriculture through understanding and enhancement of microbiomes. *New Phytol.* 230, 2129–2147. doi: 10.1111/nph.17319
- van Dorst, J., Bissett, A., Palmer, A. S., Brown, M., Snape, I., Stark, J. S., et al. (2014). Community fingerprinting in a sequencing world. *FEMS Microbiol. Ecol.* 89, 316–330. doi: 10.1111/1574-6941.12308
- Van Emon, J. M. (2016). The omics revolution in agricultural research. *J. Agric. Food Chem. Food Chem.* 64, 36–44. doi: 10.1021/acs.jafc.5b04515
- Van Goethem, M. W., Osborn, A. R., Bowen, B. P., Andeer, P. F., Swenson, T. L., Clum, A., et al. (2021). Long-read metagenomics of soil communities reveals phylum-specific secondary metabolite dynamics. *Commun. Biol.* 4:1302. doi: 10.1038/s42003-021-02809-4
- Vargas-Hernández, J. G. (2020). *Transition of Ecosystem Services Based on Urban Agro Ecology*. Hershey, Pennsylvania: IGO Global.
- Vestergaard, G., Schulz, S., Schöler, A., and Schloter, M. (2017). Making big data smart—how to use metagenomics to understand soil quality. *Biol. Fertil. Soils* 53, 479–484. doi: 10.1007/s00374-017-1191-3
- Wallenius, K., Rita, H., Simpanen, S., Mikkonen, A., and Niemi, R. M. (2010). Sample storage for soil enzyme activity and bacterial community profiles. *J. Microbiol. Methods* 81, 48–55. doi: 10.1016/j.mimet.2010.01.021
- Wang, S., Guo, Z., Wang, L., Zhang, Y., Jiang, F., Wang, X., et al. (2021). Wheat rhizosphere metagenome reveals newfound potential soil Zn-mobilizing Bacteria contributing to cultivars' variation in grain Zn concentration. *Front. Microbiol.* 12:689855. doi: 10.3389/fmicb.2021.689855
- Wang, Y., Hayatsu, M., and Fujii, T. (2012). Extraction of bacterial RNA from soil: challenges and solutions. *Microbes Environ.* 27, 111–121. doi: 10.1264/jsm2.me11304
- Watterson, W. J., Tanyeri, M., Watson, A. R., Cham, C. M., Shan, Y., Chang, E. B., et al. (2020). Droplet-based high-throughput cultivation for accurate screening of antibiotic resistant gut microbes. *elife* 9, 1–22. doi: 10.7554/eLife.56998
- Wind, L., Keenum, I., Gupta, S., Ray, P., Knowlton, K., Ponder, M., et al. (2021). Integrated metagenomic assessment of multiple pre-harvest control points on lettuce Resistomes at field-scale. *Front. Microbiol.* 12:683410. doi: 10.3389/fmicb.2021.683410
- Woese, C. R., and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. U. S. A.* 74, 5088–5090.

- Woese, C. R., Kandler, O., and Wheelis, M. L. (1990). Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc. Natl. Acad. Sci. U. S. A.* 87, 4576–4579. doi: 10.1073/pnas.87.12.4576
- Yang, X., Noyes, N. R., Doster, E., Martin, J. N., Linke, L. M., Magnuson, R. J., et al. (2016). Use of metagenomic shotgun sequencing technology to detect foodborne pathogens within the microbiome of the beef production chain. *Appl. Environ. Microbiol.* 82, 2433–2443. doi: 10.1128/AEM.00078-16
- Yang, T., Siddique, K. H. M., and Liu, K. (2020). Cropping systems in agriculture and their impact on soil health-A review. *Glob. Ecol. Conserv.* 23:e01118. doi: 10.1016/j.gecco.2020.e01118
- Yang, G., Wagg, C., Veresoglou, S. D., Hempel, S., and Rillig, M. C. (2018). How soil biota drive ecosystem stability. *Trends Plant Sci.* 23, 1057–1067. doi: 10.1016/j.tplants.2018.09.007
- Yap, M., Ercolini, D., Álvarez-Ordóñez, A., O'Toole, P. W., O'Sullivan, O., and Cotter, P. D. (2022). Next-generation food research: use of Meta-Omic approaches for characterizing microbial communities Along the food chain. *Annu. Rev. Food Sci. Technol.* 13, 361–384. doi: 10.1146/annurev-food-052720-010751
- Yun, D. Y., Kang, Y. G., Kim, E. H., Kim, M., Park, N. H., Choi, H. T., et al. (2018). Metabolomics approach for understanding geographical dependence of soybean leaf metabolome. *Food Res. Int.* 106, 842–852. doi: 10.1016/j.foodres.2018.01.061
- Zarraonaindia, I., Owens, S. M., Weisenhorn, P., West, K., Hampton-Marcell, J., Lax, S., et al. (2015). The soil microbiome influences grapevine-associated microbiota. *MBio* 6:e02527-14. doi: 10.1128/mBio.02527-14
- Zhang, M., He, L.-Y., Liu, Y.-S., Zhao, J.-L., Zhang, J.-N., Chen, J., et al. (2020). Variation of antibiotic resistome during commercial livestock manure composting. *Environ. Int.* 136:105458. doi: 10.1016/j.envint.2020.105458
- Zhang, Y. J., Hu, H. W., Chen, Q. L., Singh, B. K., Yan, H., Chen, D., et al. (2019). Transfer of antibiotic resistance from manure-amended soils to vegetable microbiomes. *Environ. Int.* 130:104912. doi: 10.1016/j.envint.2019.104912
- Zhou, A., He, Z., Qin, Y., Lu, Z., Deng, Y., Tu, Q., et al. (2013). Stress Chip as a high-throughput tool for assessing microbial community responses to environmental stresses. *Environ. Sci. Technol.* 47, 9841–9849. doi: 10.1021/es4018656

Conflict of Interest: BK was employed by the company Novasol Experts.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Djemiel, Dequiedt, Karimi, Cottin, Horrigue, Bailly, Boutaleb, Sadet-Bourgeteau, Maron, Chemidlin Prévost-Bouré, Ranjard and Terrat. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.