



**HAL**  
open science

## Interactions between proteins, phytic acid and calcium in Fava Bean concentrate

Tiffany Amat, Ali Assifaoui, Christophe Schmitt, Rémi Saurel

### ► To cite this version:

Tiffany Amat, Ali Assifaoui, Christophe Schmitt, Rémi Saurel. Interactions between proteins, phytic acid and calcium in Fava Bean concentrate. 2nd NIZO Plant Protein Functionality Conference, Oct 2022, Online, France. hal-03932226

**HAL Id: hal-03932226**

**<https://institut-agro-dijon.hal.science/hal-03932226v1>**

Submitted on 10 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## INTRODUCTION


Alternative legume proteins such as fava bean become of a huge interest for their economic and ecologic advantages as well as for their health benefits. They are rich in protein and a good source of essential nutrients and minerals. However, these advantages can be partially impaired due to their high phytic acid (PA) content. This natural polyphosphate is a major antinutrient in plant-based foods, as it can bind minerals (particularly calcium) and proteins, thereby reducing protein digestibility and subsequent mineral bioavailability. Complexes formed are insoluble and unavailable for absorption, thus lowering the nutritional value of these pulses. Molecular interactions are influenced by the environmental medium including pH, ionic strength and molar concentrations modulating the stability of these complexes during protein extraction. Protein/phytic acid/calcium complexes stability is of high relevance for food processing affecting not only structure but also functional and nutritional properties of proteins in legume-based foods. In this context, the present study investigates possible associations between fava bean proteins (FP), phytate (PA), and calcium (Ca) involved in concentrate ingredient.

## I - INGREDIENT


Table 1: Comparison of protein, phytic acid and calcium contents of fava bean concentrate powder.

	Concentrate
Processing method	Dry processing (air classification)
Protein (g/100 g)	51.7 ± 1.8
Phytic acid (PA) (g/100 g)	3.2 ± 0.7
Total calcium (mg/100 g)	127.1

## II - METHODS




## III - RESULTS



### Techniques to investigate complexes

Calcium chelation by EDTA addition


Protein removal (> 5 kDa) by Ultrafiltration



Ultrafiltration processing



Millipore Kwick Lab Cassette Pellicon® 2 Mini Ultrafiltration Biomax® Module filtration area 0.1 m<sup>2</sup>, NMWCO 5 kDa - P ≈ 2-2.5 bar

## IV - HYPOTHESIS



**pH 6.5:**

- Protein EDTA ratio  $\geq 25\%$ .
- Insoluble Binary Proteins/Ca<sup>2+</sup> complexes.
- $\rightarrow$  pH > pI ( $\approx 4-5$ ): proteins negatively charged.



**pH 7.5:**

- $\approx 40\%$  soluble PA/Protein/(Ca<sup>2+</sup>) complexes.
- PA EDTA Filtrate ratio  $\geq 8.5\%$ .
- Small % of soluble Ternary complexes.
- $\approx 90\%$  of soluble Binary PA/Protein complexes.





Figure 5: Proposed mechanisms of interactions between fava bean proteins, phytic acid and calcium depending on pH conditions.

## CONCLUSION

This study investigates possible mechanisms of interaction between phytic acid (PA), calcium (Ca) and fava bean proteins from concentrate (FPC). For the range of pH considered (6.5 to 7.5), results show the increase of the amount of soluble protein with pH raising. In addition, insoluble binary FPC/Ca complexes have been considered at pH 6.5 with no variation of phytic acid contents. The ultrafiltration process on concentrate highlights that around 40% of FPC/PA/(Ca) complexes are soluble. Even more, at pH 7.5, it should be considered a small percentage of ternary soluble complexes. Further work is needed to clarify these phenomena depending on ingredient process history. The role of protein fractions in such interaction represents an area of interest.