



HAL
open science

Spectral Estimate for Stable Signals with P-Adic Time and Optimal Selection of Smoothing Parameter

Rachid Sabre, Walid Horrigue

► **To cite this version:**

Rachid Sabre, Walid Horrigue. Spectral Estimate for Stable Signals with P-Adic Time and Optimal Selection of Smoothing Parameter. *Signal & Image Processing: An International Journal*, 2023, 14 (2/3), pp.1-9. 10.5121/sipij.2023.14301 . hal-04158343

HAL Id: hal-04158343

<https://institut-agro-dijon.hal.science/hal-04158343v1>

Submitted on 11 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

SPECTRAL ESTIMATE FOR STABLE SIGNALS WITH P-ADIC TIME AND OPTIMAL SELECTION OF SMOOTHING PARAMETER

Rachid SABRE¹ and Walid HORRIGUE²

¹Biogeosciences (UMR CNRS/uB 6282) University of Burgundy/ Institut Agro, 26, Bd
Docteur Petitjean Dijon, France

²Agroécologie, INRAE, Institut Agro, Univ. Bourgogne, Univ. Bourgogne Franche-
Comté, F-21000 Dijon, France

ABSTRACT

The spectral density of stable signals with p-adic times is already estimated under various conditions. The estimate is made by constructing a periodogram that is subsequently smoothed by a spectral window. It is clear that the convergence rate of this estimator depends on the bandwidth of the spectral window (called the smoothing parameter). This work gives a method to select the smoothing parameter in an optimal way, i.e. the estimator converges to the spectral density with the best rate.

The method is inspired by the cross-validation method, which consists in minimizing the estimate of the integrated square error.

KEYWORDS

Spectral density, p-adic processes, alpha stable, random field.

1. INTRODUCTION

This work concerns the family of alpha stable random fields, which are known by infinite variance (infinite energy). These processes are widely used models for many phenomena in several fields such as physics, biology, electronics and electricity, hydrology, economics, communications and radar applications... see [1]-[12]. The paper [13] studied the estimation of the spectral density of these processes when they are continuous time, and in [14]-[15] when the process is discrete time. The article [16] extended this work to stable random fields with p-adic time. More specifically, the process has the following spectral representation:

$$X(t_1, t_2) = \int_{Q_p^2} e^{i\langle t_1\lambda_1 + t_2\lambda_2 \rangle} dM(\lambda_1, \lambda_2);$$

$(t_1, t_2) \in Q_p^2$ where Q_p^2 is the field of p-adic numbers and M is a alpha stable random measure with a control measure m. The paper [16] studied the case where the measure m is continuous with respect to the Haar measure:

$dm(x_1, x_2) = \Phi(x_1, x_2)d\mathcal{H}(x_1, x_2)$, $(x_1, x_2) \in Q_p^2$. The density function Φ is called the spectral density of the process X. The paper [16] constructed a modified periodogram by observing the process on the ball U_n . This periodogram was then smoothed by a spectral window in order to

have an asymptotically unbiased consistent estimator of the spectral density. It is logical that the convergence rate of this estimator depends on the bandwidth of the spectral window (smoothing parameter).

The goal of this work is to propose a method for selecting the optimal smoothing parameter, allowing the best rate of convergence of the estimator. This method uses the technique of cross-validation, which has proven itself in various fields of statistics.

The choice of p-adic numbers is motivated by a great use in spatial discretization. P-adic numbers have provided solutions to many questions in physics, including questions related to string theory (related to the p-adic quantum domain) and those related to hierarchically structured fractal behaviors. They were also used in: turbulence theory, dynamical systems, statistical physics, biology, see [17]-[21]. The 2-adic number which is a special case used for computer design see [22]. Modeling in quantum mechanics used p-adic statistics [23]-[25] based on probabilistic calculations when the number of trials is infinitely large. Bernoulli's classical probabilities have been studied by [26]. The paper [27] focused on the p-adic probability theory of stochastic processes. The article [28] develops the theory of stochastic integrals with respect to p-adic Brownian motion. Using the p-adic differentiation operator, papers [29]-[32] developed the properties of the trajectories of a p-adic Wiener process.

The paper [34] set up the spectral theory and Fourier transforms of stationary p-adic processes: $X(t) t \in \mathbb{Q}_p$ where \mathbb{Q}_p is the field of p-adic numbers. He also proposed a spectral density estimator by constructing a periodogram in the same way as for real-time stationary processes. The paper [35] gives an estimator from discrete-time observation: $X(\tau_k)_{k \in \mathbb{Z}}$ where $(\tau_k)_{k \in \mathbb{Z}}$ instants of observation are taken from \mathbb{Q}_p , associated with a Poisson process.

This paper is structured as follows: section 2 gives the asymptotically unbiased and consistent estimate (smoothing of the periodogram) and the results (propositions 2.1-2.5) presented in [34]. Section 3 shows that the estimator converges in probability towards the spectral density (Proposition 2.6), gives the cross-validation criterion and Theorems 3.1-3.3 show that this criterion makes it possible to find the optimal smoothing parameter. Section 4 is reserved to the conclusion, potential applications and open research issues.

2. PERIODOGRAM AND SPECTRAL DENSITY ESTIMATION

Consider a process $X = \{X_{t_1, t_2} / (t_1, t_2) \in \mathbb{Q}_p^2\}$ where \mathbb{Q}_p^2 is the field of p-adic numbers having the following integral representation

$$X_{t_1, t_2} \int_{\mathbb{Q}_p^2} e^{i \langle t_1 \lambda_1 + t_2 \lambda_2 \rangle} dM(\lambda_1, \lambda_2) \quad (1)$$

$\forall (t_1, t_2) \in \mathbb{Q}_p^2$ where M is a symmetric α stable $S\alpha S$ random measure with independent and isotropic increments. There exists a control measure m that is defined by: $m(A \times B) = [M(A \times B), M(A \times B)]_\alpha^{1/\alpha}$.

Assume that the measure m is absolutely continuous with respect to Haar measure: $dm = \Phi(x_1, x_2) d\mathbb{H}(x_1, x_2)$ where \mathbb{H} is Haar measure.

The paper [13] gave an estimator of the density Φ , called the spectral density of the process when the process is continuous real time. In [14]-[15] the estimator is studied when the process and the

random field are discrete time. The article [16] gives the estimator of the spectral density when the process is in p-adic time: $X = \{X_{t_1, t_2} / (t_1, t_2) \in Q_p^2\}$. For that, he takes the ball $U_n = \{(x, y) \in Q_p^2; |(x, y)|_p \leq p^{-n}\}$ as the observation of the process. He considers the following periodogram: For $(\lambda_1, \lambda_2) \in Q_p^2$

$$d_n(\lambda_1, \lambda_2) = A_n \operatorname{Re} \int_{U_n} e^{-i\langle t_1 \lambda_1 + t_2 \lambda_2 \rangle} p^{-n} h(t_1 p^n, t_2 p^n) X(t_1, t_2) d\mathcal{H}((t_1, t_2)) \quad (2)$$

$$H(\lambda_1, \lambda_2) = \int_{Z_p^2} h(t_1, t_2) e^{-i\langle t_1 \lambda_1 + t_2 \lambda_2 \rangle} d\mathcal{H}(t_1, t_2)$$

$$B_\alpha = \int_{Q_p^2} |\mathcal{H}(\lambda_1, \lambda_2)|^\alpha d\mathcal{H}(\lambda_1, \lambda_2) < +\infty$$

$$H_n(\lambda_1, \lambda_2) = \left(\frac{p^{2n}}{B_\alpha}\right)^{\frac{1}{\alpha}} H(p^{-n}\lambda_1, p^{-n}\lambda_2) = A_n H(p^{-n}\lambda_1, p^{-n}\lambda_2)$$

Therefore, $A_n = \left(\frac{p^{2n}}{B_\alpha}\right)^{\frac{1}{\alpha}}$.

$$\int_{Q_p^2} |H_n(\lambda_1, \lambda_2)|^\alpha d\mathcal{H}(\lambda_1, \lambda_2) = \int_{Q_p^2} \frac{p^{2n}}{B_\alpha} |H(p^{-n}\lambda_1, p^{-n}\lambda_2)|^\alpha d\mathcal{H}(\lambda_1, \lambda_2)$$

$$= \frac{p^{2n}}{B_\alpha} p^{-2n} \int_{Q_p^2} |H(v_1, v_2)|^\alpha d\mathcal{H}(v_1, v_2)$$

Since $|p^{-n}|_p = p^{+n}$, they obtain

$$\int_{Q_p^2} |H_n(\lambda_1, \lambda_2)|^\alpha d\mathcal{H}(\lambda_1, \lambda_2) = 1$$

The following propositions 2.1-2.5 are proved in [16]

Proposition 2.1 Let

$\Psi_n(\lambda_1, \lambda_2) = \int_{Q_p^2} |H_n(\lambda_1 - u_1, \lambda_2 - u_2)|^\alpha \Phi(u_1, u_2) d\mathcal{H}(u_1, u_2)$. If Φ is a continuous and bounded function, then $B_\alpha(\Psi_n(\lambda_1, \lambda_2) - \Phi(\lambda_1, \lambda_2))$ converges to zero as n tends to infinity.

Proposition 2.2 Let $(\lambda_1, \lambda_2) \in Q_p^2$ the characteristic function of $d_n(\lambda_1, \lambda_2)$, $E \exp\{i r d_n(\lambda_1, \lambda_2)\}$, converges to $\exp\{-C_\alpha |r|^\alpha \Phi(\lambda_1, \lambda_2)\}$.

The periodogram is modified as follows:

$$I_n(\lambda_1, \lambda_2) C_{q,\alpha} |d_n(\lambda_1, \lambda_2)|^q, \quad (3)$$

where $0 < q < 2$ and the normalization constant is given by $C_{(q,\alpha)} = \frac{D_q}{F_{q,\alpha} C_\alpha^{q/\alpha}}$ where $D_q =$

$$\int \frac{1 - \cos(u)}{|u|^{1+q}} du \text{ and } F_{q,\alpha} = \int \frac{1 - \exp(-|u|^\alpha)}{|u|^{1+q}} du$$

$$C_\alpha = (\alpha\pi)^{-1} \int_0^\pi |\cos(\theta)|^\alpha d\theta,$$

Propositions 2.3 Let $(\lambda_1, \lambda_2) \in Q_p^2$, then $E I_n(\lambda_1, \lambda_2) = (\Psi_n(\lambda_1, \lambda_2))^{\frac{q}{\alpha}}$ and $I_n(\lambda_1, \lambda_2)$ is an asymptotically unbiased estimator of the spectral density but not consistent

$E I_n(\lambda_1, \lambda_2) - (\Phi(\lambda_1, \lambda_2))^{q/\alpha} = o(1)$ and $Var I_n(\lambda_1, \lambda_2) - V_{\alpha,q}(\Phi(\lambda_1, \lambda_2))^2$, with $V_{\alpha,q} = \frac{C_{2q,\alpha}^2}{C_{2q,\alpha}} - 1$.

In order to have an asymptotically and consistent estimate, we smooth the periodogram that was modified using a spectral window.

$$f_n(\lambda_1, \lambda_2) = \int_{Q_p^2} W_n(\lambda_1 - u_1, \lambda_2 - u_2) I_n(u_1, u_2) d\mathcal{H}(u_1, u_2)$$

where $W_n(x_1, x_2) = |M_n|_p W(x_1 M_n, x_2 M_n)$ such that

$$M_n \rightarrow \infty ; \frac{M_n}{n} \rightarrow 0; |M_n|_p \rightarrow 0 \text{ and } \frac{|M_n|_p}{|p^n|_p} \rightarrow \infty. \quad (4)$$

The function W is an even nonnegative function vanishing outside $[-1,1]^2$ and $\int_{Q_p^2} W(v_1, v_2) d\mathcal{H}(v_1; v_2) = 1$.

Proposition 2.4 Let $(\lambda_1, \lambda_2) \in Q_p^2$ and $Bias(f_n(\lambda_1, \lambda_2)) = E[f_n(\lambda_1, \lambda_2)] - (\Phi(\lambda_1, \lambda_2))^{p/\alpha}$, then $Bias(f_n(\lambda_1, \lambda_2)) = o(1)$. Moreover, if Φ verifies $|\Phi(x_1, x_2) - \Phi(y_1, y_2)| \leq cste |x_1 - y_1, x_2 - y_2|_p^{-k}$, then, $Bias(f_n(\lambda_1, \lambda_2)) = O\left(\frac{1}{|M_n|_p^{\frac{kp}{\alpha}}}\right)$.

Proposition 2.5 Let (λ_1, λ_2) be in Q_p^2 . Assume that $\Phi \in L^1_{Q_p^2}$. Then $Var(f_n(\lambda_1, \lambda_2)) = O(p^{-n} M_n^{-3n})$.

From propositions 2.4 and 2.5, we show in the following proposition that $(f_n(\lambda_1, \lambda_2))^{\frac{\alpha}{q}}$ converges to $\phi(\lambda_1, \lambda_2)$ in probability.

Proposition 2.6 Let λ_1, λ_2 p -adic numbers such that $\phi(\lambda_1, \lambda_2) > 0$. Then, $(f_n(\lambda_1, \lambda_2))^{\frac{\alpha}{q}}$ converges in probability to $\phi(\lambda_1, \lambda_2)$.

Proof We show that $f_n(\lambda_1, \lambda_2)$ converges in mean quadratic to $\phi(\lambda_1, \lambda_2)^{\frac{q}{\alpha}}$. $E \left| f_n(\lambda_1, \lambda_2) - \phi(\lambda_1, \lambda_2)^{\frac{q}{\alpha}} \right|^2 = E f_n(\lambda_1, \lambda_2) - \phi(\lambda_1, \lambda_2)^{\frac{q}{\alpha}})^2 + Var(f_n(\lambda_1, \lambda_2))$. Then, from proposition 2.3, $E \left| f_n(\lambda_1, \lambda_2) - \phi(\lambda_1, \lambda_2)^{\frac{q}{\alpha}} \right|^2$ converges to zero. Thus, $(f_n(\lambda_1, \lambda_2))^{\frac{\alpha}{q}}$ converges to $\phi(\lambda_1, \lambda_2)$ in probability.

It is obvious that the choice of M_n plays an important role, since the convergence rates depend on this smoothing parameter. Articles [35]-[36] use the cross-validation method to optimize the choice of parameters when the process is real-time. The objective of this work is to give a criterion for the selection of the smoothing parameters used in the estimation of the spectral density Φ . Let's note by $f(x_1, x_2) = (\Phi(x_1, x_2))^{\frac{q}{\alpha}}$ and $h = \frac{1}{M_n}$ the width of the two spectral windows. We are therefore looking for a criterion $CV(h)$ allowing us to select h to minimize the mean integrated square error (MISE), where

$$MISE(h) = \int \int E [f_n(x_1, x_2) - f(x_1, x_2)]^2 \rho(x_1, x_2) dx_1 dx_2, \quad (5)$$

ρ being a weight function that is assumed to be known and zero outside of $[0,2\pi] \times [0,2\pi]$. Although MISE(h) is a suitable measure for the squared error of the estimator, it cannot choose h, since it itself depends on the unknown function f. We adopt the cross-validation method proposed in [35]. Indeed, consider the integrated square error (ISE) defined by:

$$ISE(h) = \int \int [f_n(x_1, x_2) - f(x_1, x_2)]^2 \rho(x_1, x_2) dx_1 dx_2 = A - 2C + B$$

$$\text{where } A = \int_0^{2\pi} \int_0^{2\pi} f_n^2(x_1, x_2) \rho(x_1, x_2) dx_1 dx_2$$

$$C = \int_0^{2\pi} \int_0^{2\pi} f_n(x_1, x_2) f(x_1, x_2) \rho(x_1, x_2) dx_1 dx_2$$

$$B = \int_0^{2\pi} \int_0^{2\pi} f^2(x_1, x_2) \rho(x_1, x_2) dx_1 dx_2.$$

B being independent of h, the choice h minimizing ISE(h) amounts to choosing the h minimizing A-2C. It is clear that the term A is computable since we know f_n , whereas, in the term C, contains f unknown. We use the principle of “leave-out- I”.

3. CONSTRUCTION OF THE CROSS-VALIDATION ESTIMATOR

Let $j, j' \in \{0, 1, \dots, n-1\}$ such that $\frac{2\pi j}{n} \in U_n$ and $\frac{2\pi j'}{n} \in U_n$. The construction of “leave-out- I” consists of finding an estimator $f_n^{j,j'}(\omega_j, \omega_{j'})$ that replace $f(\omega_j, \omega_{j'})$ in the expression of C and such that $I_n(\omega_j, \omega_{j'})$ and $f_n^{j,j'}(\omega_j, \omega_{j'})$ are asymptotically independent. Thus, we can estimate C by: $\frac{1}{\bar{n}^2} \sum_{j \in A_n} \sum_{j' \in A_n} f_n^{j,j'}(\omega_j, \omega_{j'}) I_n(\omega_j, \omega_{j'}) \rho(\omega_j, \omega_{j'})$ where $\omega_j = \frac{2\pi j}{n}, \omega_{j'} = \frac{2\pi j'}{n}, \bar{n} = \lfloor \frac{n-1}{2} \rfloor$ and $A_n = \{j \in \{0, 1, \dots, n-1\} \text{ such that } \frac{2\pi j}{n} \in U_n\}$

$$f_n^{j,j'}(x_1, x_2) = \int_{U_n^2} I_n^{j,j'}(u_1, u_2) W_n(x_1 - u_1, x_2 - u_2) du_1 du_2, \text{ where}$$

$$I_n^{j,j'}(u_1, u_2) = I_n(u_1, u_2) \text{ if } (u_1, u_2) \notin B_{j,j'}$$

$$I_n^{j,j'}(u_1, u_2) = \theta_1(u_1, u_2) I_n(\omega_{j-1}, \omega_{j'-1}) +$$

$$\theta_2(u_1, u_2) I_n(\omega_{j+1}, \omega_{j'-1}) +$$

$$\theta_3(u_1, u_2) I_n(\omega_{j-1}, \omega_{j'+1}) +$$

$$\theta_4(u_1, u_2) I_n(\omega_{j+1}, \omega_{j'+1}) \quad \text{otherwise}$$

$B_{j,j'} =]\omega_{j-1}, \omega_{j+1}[\times]\omega_{j'-1}, \omega_{j'+1}[$. The construction of $I_n^{j,j'}(u_1, u_2)$ where $(u_1, u_2) \in A_{j,j'}$ is done as if I_n was bi-linear. In this case,

$$\theta_1(u_1, u_2) = \alpha\beta ; \theta_2(u_1, u_2) = (1 - \alpha)\beta ; \theta_3(u_1, u_2) = \alpha(1 - \beta) \text{ and}$$

$$\theta_4(u_1, u_2) = (1 - \alpha)(1 - \beta) \text{ where } \alpha = \frac{u_1 - \omega_{j+1}}{\omega_{j-1} - \omega_{j+1}} \text{ and } \beta = \frac{u_2 - \omega_{j'+1}}{\omega_{j'-1} - \omega_{j'+1}}.$$

The following proposition shows that $f_n^{j,j'}$ is an estimator asymptotically unbiased of the function f.

Theorem 3.1 For all $(x_1, x_2) \in Q_p^2$

$$E \left[f_n^{j,j'}(x_1, x_2) - f_n(x_1, x_2) \right] = O \left(\frac{1}{n^2} \right).$$

From this result, we establish our criterion, noted CV "cross validation" defined by:

$$CV(h) = CV_1(h) + \int_{U_n^2} f^2(u_1, u_2) \rho(u_1, u_2) du_1 du_2$$

$$\text{where } CV_1(h) = \int_{U_n^2} f_n^2(u_1, u_2) \rho(u_1, u_2) du_1 du_2 -$$

$$\frac{2}{n^2} \sum_{j \in A_n} \sum_{j' \in A_n} f_n^{j, j'}(w_j, w_{j'}) I_n(w_j, w_{j'}) \rho(w_j, w_{j'})$$

The widths of spectral windows will be chosen at the points \hat{h} minimizing the criterion $CV(h)$:

$$\hat{h}_1 = \underset{h}{\operatorname{argmin}} CV(h) = \underset{h}{\operatorname{argmin}} CV_1(h) \quad (6)$$

To facilitate writing without losing generality, we consider:

$$\rho(u_1, u_2) = \frac{1}{2\pi} \text{ on } U_n^2 \text{ and zero outside.}$$

We will show results similar to those given in [37]. This is to show that on average, the criterion $CV(h)$ and $ISE(h)$ are asymptotically close and that the variance of $CV(h)$ is asymptotically zero. Thus, the parameters \hat{h} minimizing the criterion $CV(h)$ also minimize the integral squared error (ISE) when n is large enough. These results are stated in the following theorem.

Theorem 3.2 *We have*

$$|E\{CV(h) - ISE(h)\}| = O\left(\frac{1}{n^2}\right).$$

$$\operatorname{var}\{CV(h)\} = O\left(\frac{1}{n^2 h^2}\right)$$

Thus, since

$$E\{[CV(h) - MISE(h)]^2\} =$$

$$\operatorname{var}\{CV(h)\} + [E\{CV(h) - MISE(h)\}]^2 = O\left(\frac{1}{n^2 h^2}\right).$$

The widths of the spectral windows \hat{h}_1 and \hat{h}_2 obtained by cross validation, defined in (6), are asymptotically optimal, i.e. the integrated square error at \hat{h} converges in probability to the small integrated square error.

Theorem 3.3 *The width of the spectral windows \hat{h} obtained by cross validation are asymptotically optimal:*

$$\frac{ISE(\hat{h})}{ISE(\hat{\hat{h}})} \rightarrow 1 \quad \text{in probability, where}$$

$$\hat{h} = \underset{h}{\operatorname{argmin}} CV(h) \quad \text{and} \quad (\hat{\hat{h}}) = \underset{h}{\operatorname{argmin}} ISE(h).$$

To show this result, we use the similar technique used in [38].

4. CONCLUSION

The method proposed in this work consists in giving the smoothing parameter optimizing the estimate of the spectral density for an alpha-stable process in p-adic time. To achieve this, we used the cross-validation technique, which is well suited to this kind of situation. This work can

be used in various fields of application to model phenomena whose variance is large enough such as:

- Drones take Dynamic Images. These images are often disturbed by climatic conditions. They can have a large variance and be modeled by a alpha stable random field.

- The presence of certain microorganisms in agricultural soil varies significantly can be considered as an alphastable random field.

This work will be extended to a more general case where the measure is mixed.

ACKNOWLEDGEMENTS

I would like to thank the referees for their interest in this paper and their valuable comments and suggestions. I also thank Taouakoul Brahim for proof reading.

REFERENCES

- [1] C. Gogineni, S. P. Talebi, S. Werner and D. P. Mandic, (2020), "Fractional-Order Correntropy Adaptive Filters for Distributed Processing of α -Stable Signals," in IEEE Signal Processing Letters, vol. 27, pp. 1884-1888.
- [2] X. Yan, G. Liu, H. -C. Wu, G. Zhang, Q. Wang and Y. Wu, (2020), "Robust Modulation Classification Over α -Stable Noise Using Graph-Based Fractional Lower-Order Cyclic Spectrum Analysis," in IEEE Transactions on Vehicular Technology, vol. 69, no. 3, pp. 2836-2849, 2020.
- [3] C.L. Nikias and M. Shao, (1995) "Signal Processing with Alpha-Stable Distributions and Applications", Wiley, New York.
- [4] N. Azzaoui, L. Clavier, R. Sabre, (2002), "Path delay model based on stable distribution for the 60GHz indoor channel", IEEE GLOBECOM, IEEE, pp. 441-467.
- [5] J.P. Monillet and Yu. Kegen, (2015), "Modeling Geodetic Processes with Levy alpha-Stable Distribution and FARIMA", Mathematical Geosciences. 47, no. 6, pp627-646.
- [6] I. Wahyuni, R. Sabre, (2019), "Multifocous Image Fusion Using Laplacian Pyramid Technique Based on Alpha Stable Filter" CRPASE, Vol. 05(02), 58-62.
- [7] M. Pereyra and H. Batalia, (2012), "Modeling Ultrasound Echoes in Skin Tissues Using Symmetric alpha-Stable Processes", IEEE Transactions on Ultrasonics, Ferroelectrics, and Fr 60 equen-cy Control, 59, no. 1, pp. 60-72.
- [8] Wu. Ligang and W. Zidong, 2015), " Filtering and Control for Classes of Two-Dimensional Systems", The series Studies in Systems of, Decision and Control, 18 pp. 1-29.
- [9] R. Sabre, (2020), "Wavelet Decomposition and Alpha Stable Fusion", Signal Image Processing: An International Journal (SIPIJ), Vol.11, No.1, pp. 11-24.
- [10] X. Zhong and A.B. Premkumar, (2012), "Particle filtering for acoustic source tracking in impulsive noise with alpha-stable process", IEEE Sensors Journal, 13, no. 2, pp. 589 – 600.
- [11] K. Panki, K. Takumagai and W. Jiang, "Laws of the iterated logarithm for symmetric jump processes", Bernoulli, 23, no. 4A, pp. 2330-2379, 2017.
- [12] Ga. Mang, Gang Yu, and Tao Wang,(2019), "Impulsive gear fault diagnosis using adaptive Morlet wavelet filter based on alpha-stable distribution and kurtogram." IEEE Access 7, 72283-72296.
- [13] E. Masry, S. Cambanis, (1984), " Spectral density estimation for stationary stable processes", Stochastic processes and their applications, Vol. 18, pp.1-31,.
- [14] R. Sabre, (1994), "Estimation de la densité de la mesure spectrale mixte pour un processus symétrique stable strictement stationnaire", C. R. Acad. Sci. Paris, t. 319, Série I, pp.1307-1310.
- [15] R. Sabre, (1995), "Spectral density estimation for stationary stable random fields ", Applications Mathematicae., Vol. 23, N. 2, pp. 107-133.
- [16] R. Sabre, (2019), " Mixed Spectra Estimation for Stable P-adic Random Fields", International Journal of Signal Processing Systems Vol. 7 , No. 2.
- [17] S.V. kozyrev, (2008) "Toward an ultrametric theory of turbulence", Theoretical and Mathematical Physics, Vol. 157, N. 3, pp. 1713-1722.

- [18] Li. Hua-Chieh, (2001)"On heights of p-adic dynamical systems", Proceeding of the american mathematical society, Vol.130, N. 2, pp. 379-386.
- [19] F. Ávila, (2020), "The frame of the p-adic numbers", Topology and its Applications, Vol. 273.
- [20] B. Dragovich, A.Yu. Dragovich, "A p -adic model of DNA sequence and genetic code", Mathematics and Statistics, Vol. 1, N. 1, pp. 34-41, 2009.
- [21] O. Shor, A. Glik, A. Yaniv-Rosenfeld, A. Valevski, A. Weizman, A. Khrennikov , F.Benninger, V. Ignatov, (2021)"EEG p-adic quantum potential accurately identifies depression, schizophrenia and cognitive decline", Plos one, p-Adic morphology, biomorphic structures and number theory, Biosystems, Vol. 205.
- [22] A. Klapper, M. Goresky, (1994) "2-Adic shift registers", Lecture Notes in Computer Science, 809, pp. 174-178, 1994.
- [23] R. Cianci, A. Khrennikov, " Can p-adic numbers be useful to regularize divergent expectation values of quantum observables", International Journal of Theoretical Physics, Vol. 33, N. 6, pp. 1217-1228, 1994.
- [24] D. Badziahin, Y. Bugeaud, J. Schleisitz, "On simultaneous rational approximation to a p-adic number and its integral powers", II. Proceedings of the Edinburgh Mathematical Society, Vol. 64, N. 2, pp. 317-337, 2021.
- [25] M.L.Lapidus, L. Hùng, M.van Frankenhuijsen, "p-Adic Fractal Strings of Arbitrary Rational Dimensions and Cantor Strings", P-Adic Num Ultramet Anal Appl, Vol. 13, pp. 215–230, 2021.
- [26] A. Khrennikov, "p-Adicbehaviour of Bernoulli probabilities", Statistics and Probability Letters, 37, N. 4, pp. 375-379, 1998.
- [27] A. Khrennikov, "p-Adic probability theory and its applications. The principle of statistical, stabilization of frequencies", Theoretical and Mathematical Physics, Vol. 97, N.3, pp. 1340-1348, 1993.
- [28] K. Kamizono, "Symmetric stochastic integrals with respect to p-adic Brownian motion" (2007)
- [29] V. S. Vladimirov, I. V. Volovich, and E. I. Zelenov, "p- Adic Analysis and Mathematical Physics", Fizmatlit, Moscow, 1994 [in Russian].
- [30] W.A. Zúñiga-Galindo, "Reaction-diffusion equations on complex networks and Turing patterns, via p-adic analysis", J. Math. Anal. Appl. Vol. 491, 2020.
- [31] D.Haldar, D. Singh, "p-Adic Multiwavelet Sets," P-Adic Num Ultramet Anal Appl, vol. 11, pp. 192–204, 2019.
- [32] L.Y. Hung, W. Li, Melby-Thompson, C.M. Wilson "line networks in p-adic AdS/CFT.", J. High Energ. Phys. 2019
- [33] D.R. Brillinger, "Some asymptotics of finite Fourier transforms of a stationary p-adic process", Journal of Combinatorics, Information and Systems, 16, pp. 155-169, 1991.
- [34] M. Rachdi, V. Monsan, "Asymptotic properties of p-adic spectral estimates of second order", J. of combinatoirics Information and System Sciences, Vol. 24, N. 2, pp. 113-142, 1999.
- [35] A.W. Bowman, "An alternative method of cross-validation for smoothing of density estimates", Biometrika, Vol. 71, 253-360, 1984.
- [36] C.M. Hurvich. , K.L. Beltrao, "Cross-validatory choice of a spectrum estimate and its connection with aic", J. Time Series. Analy, Vol. 11, pp. 121-137, 1990.
- [37] G. Gregoire, "Least squares cross-validation for counting process intensities", Scand. J. Statist., Vol. 20, pp.343-360, 1993.
- [38] J. Hart, P. Vieu, "Data-driven bandwidth choice for density estimation based on dependent data", Ann. Stat., Vol. 18, pp. 873-890, 1990.

AUTHORS

Rachid Sabre received the PhD degree in statistics from the University of Rouen, Rouen, France, in 1993 and Habilitation (HdR) from the University of Burgundy, Dijon, France, in 2003. He joined Agrosup Dijon, Dijon, France, in 1995, where he is an Associate Professor. From 1998 through 2010, he served as a member of Institut de Mathématiques de Bourgogne, France. He was a member of the Scientific Council AgroSup Dijon from 2009 to 2013. Since 2012, he has been a member of Laboratoire Electronique, Informatique, et Image (Le2i), France. He is author/co-author of numerous papers in scientific and technical journals and conference proceedings. His research interests lie in areas of statistical process and spectral analysis for signal and image processing.

Walid Horrigue received the PhD degree in statistics from the University of Lille, France, in 2012. He joined UMR “Agroécologie Equipe Biocom” INRA (National Institute for Agricultural Research) Dijon, France, in 2013, as a researcher where he develops research on modeling the microbiological organism in agricultural soil. He is author/co-author of some papers in scientific and technical journals and conference proceedings. His research interests lie in areas of nonparametric statistic “functional estimate with censored data”