

The Hidden Sides of Sourness Perception: Is Acid just a Basic Taste?

Carole Honoré-Chedozeau, Lucie Moyat, Méven Otheguy, Pierre Martini, Dominique Valentin

▶ To cite this version:

Carole Honoré-Chedozeau, Lucie Moyat, Méven Otheguy, Pierre Martini, Dominique Valentin. The Hidden Sides of Sourness Perception: Is Acid just a Basic Taste?. 11. European Conference on Sensory and Consumer Research: A Sense of Global Culture (EuroSense 2024), Sep 2024, Dublin, Ireland. . hal-04703925

HAL Id: hal-04703925 https://institut-agro-dijon.hal.science/hal-04703925v1

Submitted on 20 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Hidden Sides of Sourness Perception: Is Acid just a Basic Taste?



Carole HONORE-CHEDOZEAU¹, Lucie MOYAT¹, Méven OTHEGUY¹, Pierre MARTINI¹, Dominique VALENTIN²

- ¹: Pôle sensoriel IFV-SICAREX Beaujolais, 210 Boulevard Victor Vermorel, CS 60320, F-69661 Villefranche-sur-Saône Cedex, France
- ²: Centre des Sciences du Goût et de l'Alimentation, CNRS, INRAE, Institut Agro, Université de Bourgogne, F-21000 Dijon, France.




Sourness perception in sensory studies for food & beverages



Sour taste is considered as a basic taste while there is a variety of organic acids in the composition of many food and beverages products.

Different acids are also empirically used as reference standards to familiarize or train panels for descriptive profiles (e.g, lactic acid for yoghurt or tartaric acid for wine). Are organic acids sensorially different from each other?

Comparison of 4 organic acids commonly found in wine





Equi-mass concentration





3 series of 6 TETRAD tests comparing varying water solutions with 12 French wine professionals



Identification test & CATA method for the 4 acids with the best water condition after a familiarization step

Lemon
Apple

Metallic Bitterness ■ Astringency ■ Acidulous

Fruity
Freshness


Lactic Bold ☐ Roundness ☐ Other

with 110 French oenologists from all over France

Main results: a discrimination that needs to be better understood




Tetrad tests showed significant differences at alpha = 0,01% between all acids in the equimolar condition only.



- ✓ Only 15% of participants identified all the organic acids.
- ✓ Recurring inversions observed between tartaric and citric acids among oenologists (60%).
- ✓ Tactile sensations like rough, hard, aggressive and temporal descriptors were also elicited by few participants.

CA factor map on CATA data



Towards a multisensory perception of sourness?

- > Only the equimolar condition permitted to discriminate between the 4 organic acids in water solution
- > Identification test showed high variability of the percentage of correct identification with recurring inversions observed between tartaric and citric acids among oenologists.
- > Each acid was described differently with multiple sensory markers.
- > This study highlighted a discrimination between organic acids under water condition paving the way for possible sensory identification.